

MARKSCHEME

May 2000


CHEMISTRY

Higher Level

Paper 3

OPTION C – HUMAN BIOCHEMISTRY

C1. (a) (i)



(Award [1] for either circled C and [1] for the whole structure.)

[2]

(ii) In the ring structure of glucose, on the C₁ atom/the “carbonyl” C the H/OH are in different positions in α/β
OR illustration of this (diagrammatically).

[1]

[1]

(b) (i) glucose and fructose

[2]

(ii) glucose (and glucose)

[1]

(c) *(Award [1] for any of the below.)*

Food or energy reserves/resources/stores/glycogen/starch

Structure/cell walls/cellulose/chitin.

[1]

Total [8 marks]

C2. (a) 6. [1]

(b) (i) Chromatography and electrophoresis. [2]

(ii) *(Award up to [4] for the following points for EITHER paper chromatography OR electrophoresis.)*

Paper chromatography:

hydrolyse/release amino acids/heat with acid; [1]

place sample spot on paper; [1]

place paper in solvent (or suitable named solvent); [1]

compare distances travelled/ R_f values with known values. [1]

OR Electrophoresis:

hydrolysis; [1]

'loading' onto origin; [1]

variable voltage/distance moved from origin; [1]

compare isoelectric points (standards) *etc.* [1]

(c)	<u>pH 4.5</u>	<u>pH 6</u>	<u>pH 7.5</u>	
	$\text{H}_3\text{N}^+—$	$\text{H}_3\text{N}^+—\text{COO}^-$	$\text{H}_2\text{N}—\text{COO}^-$	
	[1]	[1]	[1]	[3]


Looking for functional groups only.

(In absence of other marks: three correct structures at wrong pH, award [1].)

Total [10 marks]

- C3. (a) **Substrate concentration:** activity/rate increases initially (first order); [1]
becomes constant/flattens out. [1]

A labelled correct diagram (i.e. axes labelled, correct shape) could score these two marks, for example:




Satisfactory explanation of one region of graph:

Many free active sites initially;
[active sites being occupied/becoming more saturated]. [1]

- (b) **Temperature:** increased rate initially; [1]
but then reduced markedly ($\rightarrow 0$); [1]

A labelled correct diagram (i.e. axes labelled, correct shape) could score these two marks, for example:



enzyme destroyed/denatured; [1]
since stabilising H bonds disrupted (or words to that effect). [1]

Total [7 marks]

OPTION D – ENVIRONMENTAL CHEMISTRY

	<u>Source</u>	<u>Reduction of emission</u>	
D1.	(a)	(i) Incomplete combustion of C-containing fuel/named fuel	[1]
			[1]
		(ii) Burning sulfur-containing fuel/coal	[1]
			[1]
		(iii) Reaction of gases in air/nitrogen and oxygen (at high temperature)	[1]
			[1]

* allow **catalytic converter** once only

(Award final mark for correct product from one of the above:)

- (i) Carbon dioxide;
 - (ii) Sulfur/sulfate/hydrogen sulfide; [1]
 - (iii) Nitrogen. [6]
- (b) One of SO₂ or NO_x (however described) [1]
- EITHER** SO₂ + H₂O ⇌ H₂SO₃ [1]
- OR** 2NO + 1½ O₂ + H₂O → 2HNO₃ (for example)

Total [8 marks]

- D2.**
- (a) Amount of oxygen needed to break down organic wastes; [1]
Reduced availability of oxygen/fewer living organisms. [1]
 - (b) Secondary treatment; [1]
Activated sludge process; [1]
Organic matter broken down/oxidised by bacteria. [1]
 - (c) Plant growth encouraged; [1]
Oxygen concentration reduced by plant decay. [1]
(Allow eutrophication as alternative to either of the above.)

Total [7 marks]

- D3.** (a) (i) Lethal dose *[1]*
Amount needed to kill 50 % of animals given the dose. *[1]*
- (ii) Advantage: Gives good indication of relative toxicities (of different chemicals) *[1]*
Disadvantage: does not indicate acceptable environmental level of chemical *[1]*
/does not help to make accurate assumptions re effect on humans.
- (b) Lead: Source: paints/ PbEt_4 in petrol, therefore exhaust gas/lead pipes in plumbing; *[1]*
Effect: brain damage (especially in children); *[1]*
Reducing: unleaded petrol/lead-free paints/use of copper or plastic pipes. *[1]*
- Nitrates: Source: leaching of nitrate fertilisers into rivers... *[1]*
Effect: stomach cancer/affects haemoglobin (in the young)/‘blue baby’ syndrome; *[1]*
Reducing: use less fertiliser/avoid use before rain is due. *[1]*

Total [10 marks]

OPTION E – CHEMICAL INDUSTRIES

E1. (a) Accept a temperature **range** 400–500 °C in **each** case. **[1]**

Pressure 150–500 atm (Haber) 1–2 atm (Contact) **[1]**

Catalyst iron/iron oxide Vanadium (pent/V) oxide **[1]+[1]**

(For each process, 3 correct conditions [2], 2 correct [1].)

(b) $N_2 + 3H_2 \rightleftharpoons 2NH_3$ (state symbols NOT required). **[1]**

(Don't penalise absence of reversible symbol.)

High temperature increases rate/gives greater rate of reaction **[1]**



But low yield of NH_3 **[1]**

Some comment on a compromise temperature **[1]**

(c) Raw Materials – naphtha, methane, other hydrocarbon (saturated); **[1]**

– high temperature/heat/catalyst *([1] for any one of the three.)* **[1]**

(Award [1] for any one of the following equations.)



etc.



Total [11 marks]

E2. (a) (Award [2] for any two of the following:)

‘close’ to C₂H₄ source;
close to industries needing polythene;
workforce;
away from residential areas
etc.

[2]

(b) Polar C—Cl bonds in PVC;
stronger intermolecular forces (than polythene).


[1]

[1]

(c) C₂H₃Cl + 2½ O₂ → 2CO₂ + H₂O + HCl (or doubled).

[1]

(Credit polymer equations if correct. Equations given are intentionally simplified.)



[1]

Comment on HCl being toxic or poisonous/no poisonous gases from polyethene.

[1]


(d) (Radical mechanism):

Free radical mentioned *e.g.* R• or A• or R—O—O•

[1]



[1]



[1]

equation for termination step, *e.g.* 2R• → R₂


[1]

(Detailed word descriptions of above may be awarded marks. If none of above marks are scored, [1] may be awarded for mention of initiation, propagation and termination.)

(Ionic mechanism):

(Ziegler) catalyst.

[1]



[1]



[1]

(Detailed word descriptions of above may be awarded marks.)

Total [14 marks]

OPTION F – FUELS AND ENERGY

- F1.** (a) (i) 219; [1]
86. [1]
- (ii) Mass number No change; [1]
Atomic number +1. [1]
- (b) (i) Time taken for activity to decrease by half (or words to that effect). [1]
- (ii) 11.7 days. [1]
Some working essential, e.g. 3-half lives mentioned. [1]
- (iii) $\frac{7}{8}$ or 0.875 or 87.5 %. [1]
- (iv) 12.5 % or $\frac{1}{8}$. [1]

Total [9 marks]

- F2.** (a) Zinc [1]
and graphite (accept carbon). [1]
- (b) Voltage – potential difference between electrodes; [1]
Power – total quantity of electricity available. [1]
- Voltage affected by the materials used; [1]
Power affected by the quantity of materials used. [1]

Total [6 marks]

- F3.** (a) Energy released when nucleus is synthesised from protons and neutrons/energy needed to split a nucleus into protons and neutrons. **[1]**
²²³Ra needs to become more stable. **[1]**
 This is achieved by losing mass/an α-particle. **[1]**


(b)

<u>Nature of Waste</u>	<u>Source</u>	<u>Characteristic</u>	<u>Storage</u>		
<u>Low-level waste</u>	Hospitals / checking welds / monitoring thickness of e.g. paper	Activity is low / short half-life / high volume	Stored until activity is reduced	[1]	[1]
<u>High-level waste</u>	Nuclear industry / military	Activity is high / long half-life / low volume	Making into glass / deep burial	[1]	[1] [6]
<i>(Award final mark for one extra point from list above.)</i>					[1]

Total [10 marks]

OPTION G – MODERN ANALYTICAL CHEMISTRY

G1. (a)



Light ions deflected more than heavy ions/> 1 signal obtained

OR ions(+) of different mass/charge ratio give > 1 line [1].

(General shape needed for full marks.)

[6]


(b) (i) $\left(35 \times \frac{75}{100}\right) + \left(37 \times \frac{25}{100}\right)$

[1]

= 35.50

[1]

(ii)



Both axes correctly labelled;

[1]

Three lines at 70, 72 and 74;

[1]

Heights of lines in correct order (70 > 72 > 74)

[1]

Total [11 marks]

G2. (a) $R_f = \frac{\text{distance travelled by 'solute'}}{\text{distance travelled by solvent}}$ [1]

(b) (i) Measure distance travelled by blue spot (centre) and solvent [1]
Divide one by the other [1]

(ii) Each dye has different attractions/affinities for the paper [1]
and the solvent (or words to that effect). [1]


(Solvent reference may be to solubility rather than attraction/affinity.)

(iii) Negligible attraction between the dye and paper [1]
compared with that of dye and solvent (or solubility of dye in solvent). [1]


*(In **absence** of the above award [1] for the distance moved by the dye = distance moved by the solvent.)*

Total [7 marks]

G3. (a)



[1]



[1]

[2]

(If both structures and bonding are correct but non-bonding electrons are not shown award a maximum of [1].)

(b)	<u>ONH₃</u>		<u>HONH₂</u>	
<u>Number of Peaks</u>	1	[1]	2	[1]
<u>Relative Areas</u>			1:2	[1]
<u>Reasoning</u>	All protons chemically equivalent (or words to that effect)	[1]	Protons in different chemical environment	[1]
				[5]

Total [7 marks]

OPTION H – FURTHER ORGANIC CHEMISTRY

- H1.** (a) Electrophilic addition. [1]
 Arrow from double bond to H^+ (or H of $H-Cl$) [1]
 Structure of carbocation ($CH_3-^+CH-CH_3$) [1]
 Arrow showing attack by Cl^- on central carbon of carbocation [1]
- (b) $CH_3CH_2CH_2Cl$ [1]
 Primary carbocation/ $CH_3CH_2CH_2^+$ is less stable or less likely to be formed (or secondary carbocation is more stable or more likely to be formed). [1]
 Explanation of different stabilities of carbocations (in terms of inductive effect or sharing of charge). [1]
- (c) (Substitution by) an electron-rich species (*e.g.* NH_3 ; X^-) [1]
 (lone pair)/Lewis base/Brønsted base
- (d) Arrow from $C-Cl$ bond to Cl atom [1]
 Structure of carbocation ($CH_3-^+CH-CH_3$) [1]
 Arrow showing attack by ^-OH on central carbon of carbocation [1]
OR
 Arrow from $C-Cl$ bond to Cl atom [1]
 Arrow showing attack by ^-OH on central carbon of halogenoalkane [1]
 Structure of intermediate (Cl and OH both bonded by --- to central C) [1]

Total [11 marks]

- H2.** (a) dichlorodifluoromethane (accept difluorodichloromethane) [1]
 1,1,2-trichloro,1,2,2-trifluoroethane (accept 1,1,2-trifluoro,1,2,2-trichloroethane) [1]
- (b) absorbs UV-radiation from the sun. [1]
- (c) (i) (Saturated) compounds with high bond energies. [1]
 (ii) C—Cl bond weaker than C—F [1]
 C—Cl more easily broken (than C—F). [1]
- (d) $\text{Cl}\bullet + \text{O}_3 \rightarrow \text{OCl}\bullet + \text{O}_2$ [1]
 (more correctly
 ClO•)

Total [7 marks]

- H3.** (a) Chiral carbon atom/C atom joined to 4 different groups [1]
 Two drawings showing enantiomers/chiral structures (object-mirror images). [2]
 (These may be incomplete showing only the 'chiral centre'.)
- (b) Light vibrating in one plane only. [1]
- Optically active compounds – rotate plane of polarisation of plane-polarised light. [1]
- When racemic mixture obtained [1]
 equimolar concentrations of stereoisomers affecting plane of polarisation equally
 and oppositely. [1]

Total [7 marks]
