



CHEMISTRY HIGHER LEVEL PAPER 3

1 hour 15 minutes

Candidate session number								
0	0							

#### **INSTRUCTIONS TO CANDIDATES**


- Write your session number in the boxes above.
- Do not open this examination paper until instructed to do so.
- Answer all of the questions from two of the Options in the spaces provided. You may continue your answers on answer sheets. Write your session number on each answer sheet, and attach them to this examination paper and your cover sheet using the tag provided.
- At the end of the examination, indicate the letters of the Options answered in the candidate box on your cover sheet and indicate the number of answer sheets used in the appropriate box on your cover sheet.

# Option A — Modern analytical chemistry

<b>A1.</b>	<sup>1</sup> HN	HNMR and IR spectroscopy both involve the absorption of electromagnetic radiation.										
	(a)	(i)	Identify the region of the electromagnetic spectrum used in <sup>1</sup> H NMR spectroscopy.	[1]								
		(ii)	Explain why tetramethylsilane (TMS) is used as a reference standard in <sup>1</sup> H NMR spectroscopy.	[1]								
	(b)	Iden	tify which of the following molecules absorbs IR radiation and explain your choice.	[2]								
			$H_2$ $O_2$ $HC1$									



**A2.** (a) The mass spectrum of an unknown compound,  $\mathbf{X}$ , of empirical formula  $C_2H_4O$  is shown below.




[Source: Cleapss Guides: L202 Spectra (Cleapss School Science Service), Sept 2000.]

(1)	Determine the relative molecular mass of <b>X</b> from the mass spectrum and deduce the formula of the molecular ion.	[2]
(ii)	Identify a fragment which gives rise to the peak at $m/z = 29$ .	[1]
(iii)	Comment on the absence of a peak at $m/z = 59$ .	[1]

#### (Question A2 continued)

(b) The IR spectrum of X is shown below.



[Source: http://modbo1.ibase.go.jp/sdbs/cgi-bin/cre\_index.cgi?lang=eng]

(i)	Use Table 17 of the Data Booklet to identify the bonds which correspond to the absorptions <b>A</b> and <b>B</b> .						
	A:						
(ii)	Deduce the name of the functional group present in $\mathbf{X}$ .	[1]					

#### (Question A2 continued)


(c) Typical proton chemical shift values are given in Table 18 of the Data Booklet. The <sup>1</sup>HNMR spectrum of **X** contains three peaks. Details of two of these are shown in the table below.

Peak	Chemical shift / ppm	Relative peak area	Splitting pattern
First	2.0	3	Singlet
Second	4.1	2	Quartet
Third			

(i)	Deduce a	possible	structure	for	X	that	is	consistent	with	the	mass,	IR	and	
	<sup>1</sup> HNMR sp	pectra.												[1]

[2]

- **A3.** Chromatography is used to test for the presence of illegal drugs in sport.
  - (a) A chromatogram of a concentrated urine sample from an athlete shows the presence of a banned substance known to have an  $R_f$  value of 0.75.
    - (i) Calculate the  $R_f$  values for **A** and **B**, and so deduce which of the spots corresponds to the banned substance. [2]



	$R_{\rm f}$ of ${\bf A}$ :	
	$R_{\rm f}$ of <b>B</b> :	
	Banned substance:	
(ii)	Suggest how the results could change if the experiment was repeated with a different solvent.	[1]



# (Question A3 continued)

[2]
!+
r <i>[4]</i>

# Option B — Human biochemistry

B1.	Prot	oteins are vital components of living systems.							
	(a)	State the general formula of 2-amino acids.							
	(b)	State <b>two</b> characteristic properties of 2-amino acids.	[2]						
	(c)	Using Table 19 of the Data Booklet, deduce the structural formula of <b>two</b> dipeptides that could be formed by the reaction of alanine with serine and state the other product of the reaction.	[3]						
		Other product of the reaction:							
	(d)	Explain the difference between the primary and secondary structure of proteins.	[2]						



# (Question B1 continued)

(e)	Explain how a sample of a protein can be analysed by electrophoresis.						


<b>B2.</b>	Enzymes are proteins which play an important role in the biochemical processes occurring in
	the body.

(a)	State the major function of enzymes in the human body.	[1]

(b)	Describe the mechanism of enzyme action in terms of structure.	[3]

(c) The graph below shows how the rate of an enzyme-catalysed reaction changes as the substrate concentration is increased.





(ii) Draw a line on the graph to represent the effect of adding a competitive inhibitor. [1]



# (Question B2 continued)

(d)	State and enzyme ac	the	effects	of	heav	y-metal	ions	and	temperature	increases on	[5]

# Option C — Chemistry in industry and technology

C1.	(a)		minium is extracted by the electrolysis of a molten mixture containing alumina, $Al_2O_3$ , g graphite electrodes.	
		(i)	State a half-equation for the reaction at the negative electrode (cathode).	[1]
		(ii)	Oxygen is produced at the positive electrode (anode). State the name of another gas produced at this electrode.	[1]
	(b)	(i)	State <b>two</b> properties of aluminium that make it suitable for use as an overhead electric cable.	[1]
		(ii)	Alloys of aluminium with nickel are used to make engine parts. Explain, by referring to the structure of these alloys, why they are less malleable than pure aluminium.	[2]




C2.	Poly	vinyl chloride (PVC) and polyethene are both polymers made from crude oil.	
	(a)	Explain why PVC is less flexible than polyethene.	[2]
	(b)	State how PVC can be made more flexible during its manufacture and explain the increase in flexibility on a molecular level.	[2]
	(c)	PVC can exist in isotactic and atactic forms. Draw the structure of the isotactic form showing a chain of at least six carbon atoms.	[1]

C3.	Nan	o-size	d 'test-tubes' with one open end, can be formed from carbon structures.						
	(a)	Desc	cribe these 'test-tubes' with reference to the structures of carbon allotropes.	[2]					
	(b)	(b) Carbon nanotubes can be used as catalysts.							
		(i)	Suggest <b>two</b> reasons why they are effective heterogeneous catalysts.	[2]					
		(ii)	State <b>one</b> potential concern associated with the use of carbon nanotubes.	[1]					
C4.			able nickel-cadmium batteries are used in portable electrical equipment and y lighting.						
	The	disch	arge process can be summarized by the equation below.						
			$2\text{NiO(OH)}(s) + \text{Cd}(s) + 2\text{H}_2\text{O}(l) \rightleftharpoons 2\text{Ni(OH)}_2(s) + \text{Cd(OH)}_2(s)$						
	(a)		e the change in oxidation number of the cadmium and deduce if it is acting as the tive or negative electrode during the discharge process.	[2]					
	(b)		tify a physical property of Cd(OH) <sub>2</sub> which allows this process to be reversed and the ery recharged.	[1]					



**C5.** The structure of 4-pentyl-4-cyanobiphenyl, a commercially available nematic crystalline material used in electrical display devices, is shown below.



(a)	Explain how the three different parts of the molecule contribute to the properties of the compound used in electrical display devices.	[3]
	CN:	
	$C_5H_{11}$ :	
(b)	Describe and explain in molecular terms the workings of a twisted nematic liquid crystal.	[4]

#### Option D — Medicines and drugs

<b>D1.</b> The discovery of penicillin was one of the most significant scientific d	liscoveries of the last centu	ry.
---	-------------------------------	-----

(a)	Describe the mode of action of penicillins in treating infectious diseases.	[2]

(b) Ampicillin is a semi-synthetic penicillin used to treat lung infections. The structure of the antibiotic is shown below.

(i)	Explain why it is important to continue to develop semi-synthetic penicillins.	[2]
(ii)	Describe how computers are used in drug design.	[2]



[2]

(Question D1 continued)

(c)	State the type of hybridization of each of the carbon atoms (I, II, and III) in the $\beta$ -lactam
	ring of ampicillin by completing the table below, and explain why the amide group is
	highly reactive.

Carbon atom	I	II	III
Hybridization			

•	٠	•	•	•	٠	•	•	•	•	 •	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	٠	•	•	٠	•	•	•	•	٠		•	٠	•	•	٠	•	•	٠		•	•	•	•	•	٠	•	٠	٠	٠		•	•	•	•	•	•	٠		•	•	•	•	•	٠	

<b>D2.</b>	(a)	Expl	ain the meaning of the terms:	
		(i)	side-effect	[1]
		(ii)	therapeutic window	[1]
		(iii)	placebo effect.	[1]
	(b)	The	effectiveness of a drug depends on the method of administration.	
		(i)	One method of injecting drugs into the body results in the drug having a very rapid effect. State the method and explain its rapid action.	[2]
		(ii)	List the <b>two</b> other methods which can be used to inject drugs into the body.	[1]
		(iii)	Identify the method of administration used to treat respiratory diseases such as asthma.	[1]




D3.	Amp	phetamine and methamphetamine are widely abused drugs.	
	(a)	State <b>one</b> short-term effect of amphetamine on the human body.	[1]
	(b)	Explain why amphetamine is classified as a <i>sympathomimetic drug</i> and relate this to its structure.	[2]
	(c)	Regular use of amphetamine and methamphetamine can lead to <i>tolerance</i> . Explain why this is potentially dangerous.	[2]
	(d)	Amphetamine exists as optical isomers. Describe how chiral auxiliaries can be used to synthesize only the desired enantiomeric form of a drug from a non-chiral starting compound.	[3]
	(e)	The structures of morphine and heroin are shown in Table 20 of the Data Booklet. Explain the increased potency of heroin compared to morphine.	[2]

# $Option \ E -- Environmental \ chemistry$

E1.		now widely accepted that the increased production of carbon dioxide is leading to al warming.	
	(a)	Describe how carbon dioxide acts as a greenhouse gas.	[2]
	(b)	Discuss the influence of increasing amounts of greenhouse gases on the environment.	[3]
E2.	The	biochemical oxygen demand (BOD) is a measure of water pollution.	
	(a)	Identify the stage of sewage treatment which removes the substances responsible for high BOD values and explain how this is done.	[2]
	(b)	Describe how the addition of nitrates or phosphates to water can increase the BOD value of a water sample.	[2]



**E3.** The concentration of some pollutants in a city was measured over a 24 hour period. The results are shown below.



(a)	Outline the cause of the increase in NO levels between 5 and 8 am.										

Outline the cause of the increase in concentration of peroxyacylnitrates (PANs) between 8 am and 2 pm.	[2]

(c)	State why a temperature inversion often results in the formation of smog.										

(d) Show, by stating an equation, how **one** of the above pollutants reacts with HO• radicals to produce a constituent of acid rain. [1]

(b)

E4.		dfill sites are used to dispose of about 90% of the world's domestic waste, but incineration ing increasingly used in some countries.	
	(a)	State <b>one</b> advantage of each method.	[2]
		Landfill:	
		Incineration:	
	(b)	Suggest why some biodegradable plastics do not decompose in landfill sites.	[1]
	(c)	High-level and low-level wastes are two types of radioactive waste. Compare the half-lives and the methods of disposal of the two types of waste.	[3]

E5.	can 1	pounds of heavy metals are one type of toxic substance found in water. Lead(II) ions, $Pb^{2\tau}$ , be removed by bubbling hydrogen sulfide, $H_2S$ , through polluted water. The solubility act of lead sulfide is $1.25 \times 10^{-28} \text{mol}^2 \text{dm}^{-6}$ at $25^{\circ}C$ .	
	(a)	Calculate the concentration of Pb <sup>2+</sup> ions in a saturated solution of lead sulfide.	[2]
	(b)	Explain how the addition of hydrogen sulfide decreases the concentration of Pb <sup>2+</sup> ions in a saturated solution.	[2]

# Option F — Food chemistry

(a)	State	e one function of a nutrient.	[
(b)	Desc	eribe the structural composition of the following nutrients:	
	(i)	fats and oils	I
	(ii)	monosaccharides.	
(c)	fish (	r is a source of arachidonic acid, $CH_3(CH_2)_4(CH=CHCH_2)_4(CH_2)_2COOH$ , and oils are a source of linolenic acid. With reference to the structure of linolenic acid in e 22 of the Data Booklet, explain why arachidonic acid has a much lower melting t compared to linolenic acid, even though it contains two more carbon atoms.	



F2.	Antioxidants	can be us	ed to pro	olong the s	helf life of	food.

	(a)	Define	the	terms
۱	a		uic	willis.

(i)

(ii)

shelf life	[1]
antioxidant.	[1]

(b) Using Table 22 of the Data Booklet, and the structure of propyl gallate (PG) below, compare the structural features of the three common antioxidants 3-BHA, BHT and PG. [4]

•	•																																			 	•	•	•
					•	•			•	•			•	•			•	•	•		 •	•		 			•					•	•	 •		 	•	•	•
																 								 						-						 			
																																							•
•	•			٠		•		 ٠	•			 ٠		•		 		•	•		 ٠	•	•	 	•			 •	•	•	 •	•	•	 ٠	•	 	٠	•	•

#### (Question F2 continued)

Long-term health benefit:

(c)	State <b>one</b> example of a common naturally occurring antioxidant and state <b>one</b> possible long-term health benefit of consuming food in which it is present.	[2]
	Antioxidant:	

.....

(d) (EDTA)<sup>4-</sup>, the ethylenediaminetetraacetate anion, is a chelate ligand with the following structure.

$$\begin{array}{c} : O: \\ \vdots \vdots \\ CH_2 \\ CH$$

It has been found to inhibit the Fe<sup>2+</sup> catalysed oxidation of raw beef. Explain why (EDTA)<sup>4-</sup> can be described as a chelate ligand. [1]

**F3.** Different conventions are used to classify enantiomers. Orange and lemon peel each contain different enantiomers of the compound limonene. One of the enantiomers is represented below.

(a) Identify the chiral centre in this enantiomer with an asterisk, \*. [1]
(b) The (+)*d*-enantiomer has the characteristic smell of oranges and the (-)*l*-enantiomer has the characteristic smell of lemons. Explain the meaning of the (+)*d* and (-)*l* symbols used in this notation. [1]
(c) The R, S notation is also used. The (+)*d*-enantiomer is often described as R-limonene and the (-)*l*-enantiomer as S-limonene. Explain what is meant by the R, S notation and state whether the structure shown is R or S. [2]

F4.	The chemical process that makes butter rancid is called oxidative rancidity. Describe, using equations, the steps involved in the free-radical chain mechanism that occurs during oxidative rancidity.	[4]
	Initiation:	
	Propagation:	
	Termination:	



#### **Option G** — Further organic chemistry

- **G1.** The carbonyl chemistry of both aldehydes and ketones involves a number of different types of reactions.
  - (a) State the formulas of the products, A–C, formed in the following reactions.

(i) 
$$(CH3)2CO \xrightarrow{CH3CH2MgBr} A + B$$
 [2]

**A**:

**B**:

(ii)  $CH_3CHO + H_2N \longrightarrow NH \longrightarrow C + H_2O$  [1]

C: .....

(b) State the type of reaction involved in the formation of C and  $H_2O$  in (a) (ii). [1]

.....

<b>G2.</b>	Etha	noic acid and chloroethanoic acid both contain the carboxyl group.	
	(a)	Explain why chloroethanoic acid is a stronger acid than ethanoic acid.	[2]
	(b)	Starting with chloromethane, outline <b>one</b> possible reaction pathway to synthesize ethanoic acid. Your answer should include:  • the reagents used	
		• a relevant chemical equation for each step.	[2]



<b>G3</b> .	(a)	Alkenes commonly react by electrophilic addition reactions. Describe the mechanism of	
		the following reaction, using curly arrows to show the movement of electron pairs, and	
		suggest the reason for the formation of the major organic product, <b>D</b> .	[5]


$$CH_2=CH(CH_2CH_3) + HI \rightarrow \mathbf{D}$$




#### (Question G3 continued)

(b) Both benzene and methylbenzene undergo electrophilic substitution reactions. State the formulas of the reagents, **E-G**, **including any catalysts used**, in the following conversions.

[3]



E: ....



F: .....

G:



# (Question G3 continued)

(c)	State	e the formulas of the products <b>H</b> and <b>I</b> , formed in the following reaction.	[2]
		$C_6H_5CH_2Br + NaOH \rightarrow H + I$	
	H:		
	I:		
(d)	(i)	State whether –OH can be described as an activating or deactivating group with respect to aromatic electrophilic substitution reactions.	[1]
	(ii)	Describe the directing effect of –OH on a benzene ring in an aromatic electrophilic substitution reaction.	[1]
	(iii)	Explain the decreased reactivity due to the presence of the $-\mathrm{NO}_2$ group in an aromatic electrophilic substitution reaction.	[2]
(e)	Cons	sider the two amines methylamine, CH <sub>3</sub> NH <sub>2</sub> , and dimethylamine, (CH <sub>3</sub> ) <sub>2</sub> NH.	
	(i)	Using Table 15 of the Data Booklet, state which of the amines is the more basic.	[1]
	(ii)	Explain the relative basicities of the two amines.	[2]

