



# INTERNATIONAL BACCALAUREATE

## MATHEMATICS

Higher Level

Tuesday 7 May 1996 (morning)

Paper 2

2 hours 30 minutes

This examination paper consists of 2 sections, Section A and Section B.

Section A consists of 4 questions.

Section B consists of 4 questions.

The maximum mark for Section A is 80.

The maximum mark for each question in Section B is 40.

The maximum mark for this paper is 120.

This examination paper consists of 11 pages.

### INSTRUCTIONS TO CANDIDATES

**DO NOT open this examination paper until instructed to do so.**

**Answer all FOUR questions from Section A and ONE question from Section B.**

**Unless otherwise stated in the question, all numerical answers must be given exactly or to three significant figures as appropriate.**

### EXAMINATION MATERIALS

#### Required/Essential:

IB Statistical Tables  
Millimetre square graph paper  
Electronic calculator  
Ruler and compasses

#### Allowed/Optional:

A simple translating dictionary for candidates not working in their own language

**FORMULAE**

**Trigonometrical identities:**

$$\sin(\alpha + \beta) = \sin \alpha \cos \beta + \cos \alpha \sin \beta$$

$$\cos(\alpha + \beta) = \cos \alpha \cos \beta - \sin \alpha \sin \beta$$

$$\tan(\alpha + \beta) = \frac{\tan \alpha + \tan \beta}{1 - \tan \alpha \tan \beta}$$

$$\sin \alpha + \sin \beta = 2 \sin \frac{\alpha + \beta}{2} \cos \frac{\alpha - \beta}{2}$$

$$\sin \alpha - \sin \beta = 2 \cos \frac{\alpha + \beta}{2} \sin \frac{\alpha - \beta}{2}$$

$$\cos \alpha + \cos \beta = 2 \cos \frac{\alpha + \beta}{2} \cos \frac{\alpha - \beta}{2}$$

$$\cos \alpha - \cos \beta = 2 \sin \frac{\alpha + \beta}{2} \sin \frac{\beta - \alpha}{2}$$

$$\cos 2\theta = 2 \cos^2 \theta - 1 = 1 - 2 \sin^2 \theta = \cos^2 \theta - \sin^2 \theta$$

$$\text{If } \tan \frac{\theta}{2} = t \text{ then } \sin \theta = \frac{2t}{1+t^2} \text{ and } \cos \theta = \frac{1-t^2}{1+t^2}$$

**Integration by parts:**

$$\int u \frac{dv}{dx} dx = uv - \int v \frac{du}{dx} dx$$

**Standard integrals:**

$$\int \frac{dx}{x^2 + a^2} = \frac{1}{a} \arctan \frac{x}{a} + c$$

$$\int \frac{dx}{\sqrt{a^2 - x^2}} = \arcsin \frac{x}{a} + c \quad (|x| < a)$$

**Statistics:**

If  $(x_1, x_2, \dots, x_n)$  occur with frequencies  $(f_1, f_2, \dots, f_n)$  then the mean  $m$  and standard deviation  $s$  are given by

$$m = \frac{\sum f_i x_i}{\sum f_i} \quad s = \sqrt{\frac{\sum f_i (x_i - m)^2}{\sum f_i}}, \quad i = 1, 2, \dots, n$$

**Binomial distribution:**

$$p_x = \binom{n}{x} p^x (1-p)^{n-x}, \quad x = 0, 1, 2, \dots, n$$

A correct answer with **no** indication of the method used will normally receive **no** marks. You are therefore advised to show your working.

### SECTION A

Answer all **FOUR** questions from this section.

1. [Maximum mark: 20]

- (i) The line  $L$  passes through the point  $P(2, 3, 1)$  and has direction  $\vec{u} = \vec{i} + 2\vec{j} - 3\vec{k}$ . A second line  $M$  passes through the point  $Q(4, 2, 0)$  and has direction  $\vec{v} = 3\vec{i} - \vec{j} + \vec{k}$ .

(a) Write down, in parametric form and using  $t$  as the parameter, the equation of the line  $M$ .

(b) Find the vector  $\vec{w} = \vec{RP}$ , where  $R$  is any point on the line  $M$ .

(c) Write down the vector  $\vec{u} \times \vec{w}$  and hence express  $|\vec{u} \times \vec{w}|$  in terms of  $t$ .

(d) Deduce that  $|\vec{u} \times \vec{w}|$  is minimised when  $t = -\frac{3}{5}$  and find this minimum value.

[12 marks]

- (ii) Write down, in polar form, the cube roots of  $-i$ . Hence, or otherwise, solve the equation

$$[(1 - i)z]^3 + i = 0,$$

expressing your answers in algebraic form (that is, in the form  $a + ib$ ).

[8 marks]

2. [Maximum mark: 22]

(a) Let the function  $x(t)$  be defined by

$$x(t) = e^{-\lambda t} \sin (pt + \alpha),$$

where  $\lambda$ ,  $p$  and  $\alpha$  are constants. Show that

$$\frac{d^2x}{dt^2} + 2\lambda \frac{dx}{dt} + (\lambda^2 + p^2)x = 0.$$

[6 marks]

(b) Given that  $\frac{dx}{dt} = -2p$  and  $\frac{d^2x}{dt^2} = -3p$ , when  $\alpha = 0$  and  $t = \frac{\pi}{p}$ , calculate the value of  $\lambda$  and show that

$$p = \frac{3\pi}{4 \ln 2}.$$

[4 marks]

(c) Now consider the different case when  $\frac{dx}{dt} = 0$  at  $t = 0$ , and  $\alpha$  is not specified. Show that

(i)  $\lambda = p \cot \alpha$ ;

(ii) the values of  $t$  when  $\frac{dx}{dt} = 0$  are in arithmetic progression, and find the common difference;

(iii) the values of  $x$  when  $\frac{dx}{dt} = 0$  are in geometric progression, and find the common ratio in terms of  $\alpha$ .

[12 marks]

3. [Maximum mark: 20]

(i) Find the area enclosed by the curve  $y = x^2 \sin x$  and the  $x$ -axis for  $0 \leq x \leq 2\pi$ , giving your answer exactly in terms of  $\pi$ . [10 marks]

(ii) (a) Derive an equation that  $a$ ,  $b$  and  $c$  must satisfy for the system of equations

$$\begin{aligned} -3x + y + 2z &= a \\ -11x + 2y + 6z &= b \\ 7x + y - 2z &= c \end{aligned}$$

to have a solution.

(b) Derive a solution when  $a = 2$  and  $b = 7$ . Is the solution unique? Explain your answer clearly. [10 marks]

4. [Maximum mark: 18]

Note: In this question all answers must be given exactly as rational numbers.

- (a) A man can invest in at most one of two companies,  $A$  and  $B$ . The probability that he invests in  $A$  is  $\frac{3}{7}$  and the probability that he invests in  $B$  is  $\frac{2}{7}$ , otherwise he makes no investment.

The probability that an investment yields a dividend is  $\frac{1}{2}$  for company  $A$  and  $\frac{2}{3}$  for company  $B$ .

The performances of the two companies are totally unrelated.

Draw a probability tree to illustrate the various outcomes and their probabilities. What is the probability that the investor receives a dividend and, given that he does, what is the probability that it was from his investment in company  $A$ ?

[8 marks]

- (b) Suppose that a woman must decide whether or not to invest in each company. The decisions she makes for each company are independent and the probability of her investing in company  $A$  is  $\frac{3}{10}$  while the probability of her investing in company  $B$  is  $\frac{6}{10}$ . Assume that there are the same probabilities of the investments yielding a dividend as in (a).

- (i) Draw a probability tree to illustrate the investment choices and whether or not a dividend is received. Include the probabilities for the various outcomes on your tree.
- (ii) If she decides to invest in both companies, what is the probability that she receives a dividend from at least one of her investments?
- (iii) What is the probability that she decides not to invest in either company?
- (iv) If she does not receive a dividend at all, what is the probability that she made no investment?

[10 marks]

**SECTION B**

Answer ONE question from this section.

**Abstract Algebra**

5. [Maximum mark: 40]

- (i) Show that  $\mathbb{R}$ , the set of real numbers, forms a group under the operation  $*$  where

$$a * b = \sqrt[3]{a^3 + b^3}$$

for all  $a, b$  in  $\mathbb{R}$ .

[11 marks]

- (ii) Consider the three points  $(2, 0)$ ,  $(1, \sqrt{3})$  and  $(-1, \sqrt{3})$  in the  $x$ - $y$  plane.

Let  $R_\theta$  denote the anticlockwise rotation about the origin through the angle  $\theta$ ,  $0 \leq \theta < 2\pi$ .

Let  $S$  be the set of anticlockwise rotations about the origin which map any one of the these points either onto itself or onto one of the other points.

- (a) List the elements of the set  $S$ .
- (b) Show that  $S$  is not a group under the composition of rotations.
- (c) Find a rotation  $R_\alpha$ , such that  $S \cup R_\alpha$  does form a group under the composition of rotations. Show that this group is cyclic and find a generator for it. Write down all of the proper subgroups of this group.

[14 marks]

- (iii) (a) Explain what is meant by stating that two groups are **isomorphic**.

(b) Consider two isomorphic groups  $\{G_1, *\}$  and  $\{G_2, \circ\}$  with identity elements  $e_1$  in  $G_1$  and  $e_2$  in  $G_2$ . Let  $f: G_1 \rightarrow G_2$  be an isomorphism,  $a_1$  any element of  $G_1$  and write  $a_2 = f(a_1)$ . Prove that

1.  $e_2 = f(e_1)$ ;


2.  $a_2^{-1} = f(a_1^{-1})$ .

[15 marks]

**Graphs and Trees**

6. [Maximum mark: 40]

(i) Consider the five graphs below



(a) By copying, and then completing the table below, write down the degrees of the vertices  $A, B, C, D$  and  $E$  in each graph and state whether the graph is Eulerian or not, and whether it is Hamiltonian or not. Whenever your answer is yes, give a circuit that verifies your answer.

	Graph I	Graph II	Graph III	Graph IV	Graph V
Degrees	4, 4, 4, 4, 4				
Eulerian?	Yes, $ABCD$ $EACEBDA$				
Hamiltonian?	Yes, $ABCDE$				

(b) Which of the following statements are true? (No proofs or counter examples are required.)

- A graph is Hamiltonian  $\Rightarrow$  it is Eulerian.
- A graph is Eulerian  $\Rightarrow$  it is Hamiltonian.
- A graph is not Hamiltonian  $\Rightarrow$  it is Eulerian.
- A graph is not Eulerian  $\Rightarrow$  it is Hamiltonian.

(c) Write down a necessary and sufficient condition for a connected graph to be Eulerian and explain why your condition is necessary.


[16 marks]

(This question continues on the following page)



(Question 6 continued)

- (ii) Use Dijkstra's Algorithm to find the shortest path, and its length, from  $A$  to  $H$  in the directed graph below, showing your working clearly in tabular fashion.



What is the length of the shortest path from  $A$  to  $G$ ?

[12 marks]

- (iii) Let  $G$  be a connected planar graph that is not a tree which has  $v$  vertices,  $e$  edges and divides the plane into  $f$  faces or regions. State Euler's formula connecting  $v$ ,  $e$  and  $f$  and verify the formula when  $e = 0$ .

Hence, assuming that the formula is true when there are  $n - 1$  edges, use mathematical induction to prove that the formula is true.

[12 marks]

Statistics

7. [Maximum mark: 40]

- (i) The data in the table below shows the number of telephone calls received per minute at a business office over a forty minute period.

Calls per minute	0	1	2	3	4	5	>5
Frequency	1	8	12	7	8	4	0

Perform a  $\chi^2$  test at the 5% level of significance to determine whether or not the above data can be modelled by a Poisson distribution with mean given by the sample mean.

[12 marks]

- (ii) Systolic blood pressures were recorded for a sample of 25 diabetic males in a given age group. The sample mean and standard deviation were 147.60 and 13.92 respectively. From a sample of 30 non-diabetic males in the same age group the corresponding values were 139.61 and 12.59 respectively. (The units are mm Hg.)

- (a) Explain in detail how to obtain a 95% confidence interval for the difference in mean blood pressure in diabetic and non-diabetic males in the population from which the samples were drawn. The distributional results and assumptions required for the derivation of such an interval should be carefully stated.

Given that  $t_{0.025} = 2.01$  for a  $t$  distribution with 53 degrees of freedom, calculate the end points of the interval. Carry out a test of the null hypothesis of equal population means for diabetic and non-diabetic males against a two sided alternative hypothesis, using a 5% significance level.

[15 marks]

- (b) In a separate study it was found that 43 out of 200 diabetic males and 41 out of 300 non-diabetic males were classified as having high blood pressure. Use a  $\chi^2$  test to investigate independence of the two classifications **diabetic** and **high blood pressure**.

[Full details of your working should be included.]

[13 marks]

**Analysis and Approximation**

8. [Maximum mark: 40]

- (i) (a) The curves of the functions  $y = x^3 - 21$  and  $y = x^2 + 2$  intersect at the point  $(x^*, y^*)$ . Find the integer  $m$  such that  $m < x^* < m + 1$ .
- (b) The value  $x^*$  satisfies both of the two equations

$$x = \sqrt[3]{x^2 + 23}; \quad x = \sqrt{x^3 - 23}.$$

Which, if any, of these two can be used as an iterative scheme of the form  $x_{n+1} = g(x_n)$ , with  $x_0 = m$ , to approximate the value  $x^*$ ? Explain clearly your decision about each scheme.

Find  $x^*$  to 3 decimal places.

[18 marks]

- (ii) (a) For  $x > 0$  and  $n \in \mathbb{N}$ , the exponential function  $e^x$  can be written as

$$e^x = 1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} \dots + \frac{x^n}{n!} + \frac{x^{n+1}}{(n+1)!} e^\theta$$

where  $0 < \theta < x$ .

Deduce that  $e^x > \frac{x^n}{n!}$  for any  $n \in \mathbb{N}$ .

- (b) By considering the case when  $n = 1$ , deduce that  $\ln t < \frac{t^p}{p}$  for  $t > 1$  and  $p \in \mathbb{R}, p > 0$ .

- (c) Deduce that

$$\ln k < 4k^{-\frac{1}{4}}, \quad k = 1, 2, 3, 4, 5, \dots$$

and hence use the comparison test to determine whether or not the series

$$\sum_{n=1}^{\infty} \frac{\ln n}{1 + n^{\frac{3}{2}}}$$

converges. You may state, without proof, the condition for the series

$$\sum_{n=1}^{\infty} \frac{1}{n^p} \text{ to converge.}$$

[22 marks]