



**MATHEMATICS
HIGHER LEVEL
PAPER 2**

Thursday 4 May 2000 (morning)

3 hours

INSTRUCTIONS TO CANDIDATES

- Do not open this examination paper until instructed to do so.
- Answer all five questions from Section A and one question from Section B.
- Unless otherwise stated in the question, all numerical answers must be given exactly or to three significant figures, as appropriate.
- Write the make and model of your calculator on the front cover of your answer booklets *e.g.* Casio *fx-7400G*, Sharp EL-9400, Texas Instruments TI-80.


You are advised to start each new question on a new page. A correct answer with **no** indication of the method used will usually receive **no** marks. You are therefore advised to show your working. In particular, where graphs from a graphic display calculator are being used to find solutions, you should sketch these graphs as part of your answer.

SECTION A

Answer all **five** questions from this section.

1. [Maximum mark: 10]

The area of the triangle shown below is 2.21 cm^2 . The length of the shortest side is $x \text{ cm}$ and the other two sides are $3x \text{ cm}$ and $(x + 3) \text{ cm}$.



- (a) Using the formula for the area of the triangle, write down an expression for $\sin \theta$ in terms of x . [2 marks]
- (b) Using the cosine rule, write down and simplify an expression for $\cos \theta$ in terms of x . [2 marks]
- (c) (i) Using your answers to parts (a) and (b), show that,


$$\left(\frac{3x^2 - 2x - 3}{2x^2} \right)^2 = 1 - \left(\frac{4.42}{3x^2} \right)^2 \quad [1 \text{ mark}]$$

(ii) Hence find

- (a) the possible values of x ; [2 marks]
- (b) the corresponding values of θ , **in radians**, using your answer to part (b) above. [3 marks]

2. [Maximum mark: 14]

The diagram below shows the graphs of $y = -x^3 + 3x^2$ and $y = g(x)$, where $g(x)$ is a polynomial of degree 3.



- (a) If $g(-2) = 0$, $g(0) = -4$, $g'(-2) = 0$, and $g'(0) = 0$ show that $g(x) = x^3 + 3x^2 - 4$. [6 marks]

The graph of $y = -x^3 + 3x^2$ is reflected in the y -axis, then translated using the vector $\begin{pmatrix} -1 \\ -1 \end{pmatrix}$ to give the graph of $y = h(x)$.

- (b) Write $h(x)$ in the form $h(x) = ax^3 + bx^2 + cx + d$. [5 marks]

The graph of $y = x^3 + 3x^2 - 4$ is obtained by applying a composition of two transformations to the graph of $y = -x^3 + 3x^2$.

- (c) State the two transformations whose composition maps the graph of $y = -x^3 + 3x^2$ onto the graph of $y = x^3 + 3x^2 - 4$ and also maps point A onto point A'. [3 marks]

3. [Maximum mark: 16]

Let $f(x) = \ln |x^5 - 3x^2|$, $-0.5 < x < 2$, $x \neq a$, $x \neq b$; (a , b are values of x for which $f(x)$ is not defined).

- (a) (i) Sketch the graph of $f(x)$, indicating on your sketch the number of zeros of $f(x)$. Show also the position of any asymptotes. [2 marks]
- (ii) Find all the zeros of $f(x)$, (that is, solve $f(x) = 0$). [3 marks]
- (b) Find the **exact** values of a and b . [3 marks]
- (c) Find $f'(x)$, and indicate clearly where $f'(x)$ is not defined. [3 marks]
- (d) Find the **exact** value of the x -coordinate of the local maximum of $f(x)$, for $0 < x < 1.5$. (You may assume that there is no point of inflexion.) [3 marks]
- (e) **Write down** the definite integral that represents the area of the region enclosed by $f(x)$ and the x -axis. (Do **not** evaluate the integral.) [2 marks]

4. [Maximum mark: 12]

A machine is set to produce bags of salt, whose weights are distributed normally, with a mean of 110 g and standard deviation of 1.142 g. If the weight of a bag of salt is less than 108 g, the bag is rejected. With these settings, 4% of the bags are rejected.

The settings of the machine are altered and it is found that 7% of the bags are rejected.

- (a) (i) If the mean has not changed, find the new standard deviation, **correct to three decimal places**. [4 marks]
- The machine is adjusted to operate with this new value of the standard deviation.
- (ii) Find the value, **correct to two decimal places**, at which the mean should be set so that only 4% of the bags are rejected. [4 marks]
- (b) With the new settings from part (a), it is found that 80% of the bags of salt have a weight which lies between A g and B g, where A and B are symmetric about the mean. Find the values of A and B , giving your answers **correct to two decimal places**. [4 marks]

5. [Maximum mark: 18]

(i) Differentiate from first principles $f(x) = \cos x$. [8 marks]

(ii) Prove by mathematical induction that $\frac{d}{dx}(x^n) = nx^{n-1}$, for all positive integer values of n . [10 marks]

SECTION B

Answer one question from this section.

Statistics

6. [Maximum mark: 30]

A computer manufacturing company buys large quantities of hard discs from several suppliers. Hard disc quality is checked by a process called RTT which gives results on a continuous scale from 0 to 100 . Based on previous experience the company assumes that the results are normally distributed with a mean of 68 and standard deviation of 3 .

Shipments arrive from suppliers on a daily basis. A sample of 10 hard discs is taken from each shipment at random and tested. If the mean of the sample is more than 67 , the shipment is accepted, otherwise it is rejected.

- (a) What is the probability that a hard disc selected at random has a result less than 67 ? [2 marks]

- (b) Find the probability that a shipment is rejected. [4 marks]

- (c) There is a \$1000 penalty each day that a shipment is rejected. A particular supplier’s hard discs have a mean of 67.5 and a standard deviation of 2.8 .
 - (i) What is the probability that this supplier’s shipment is accepted? [3 marks]

 - (ii) What is the expected penalty per 6-day week for this supplier? [6 marks]

- (d) The company’s own production of hard discs has a mean of 68 and a standard deviation of 3 . However, to keep the production within the acceptable limits, the company samples 10 hard discs every hour and examines whether the sample is accepted or rejected. During a particular hour, the following results were recorded for a sample that was randomly chosen for testing:

68.1747, 68.0473, 66.3189, 66.2735, 66.957, 66.9738,
66.1438, 67.0744, 66.1875, 67.8804

At the 5% level of significance determine whether the sample meets the company’s standard for acceptance. [7 marks]

(This question continues on the following page.)

(Question 6 continued)

- (e) Every week, the company randomly selects the test results of 1000 hard discs and checks if these results come from a normal distribution (with a mean of 68 and standard deviation of 3) or not. The following table gives the results, R , for one such test.

Results	Frequency
$56 \leq R < 59$	5
$59 \leq R < 62$	17
$62 \leq R < 65$	146
$65 \leq R < 68$	333
$68 \leq R < 71$	360
$71 \leq R < 74$	113
$R \geq 74$	26

Check, at the 5% level of significance, whether the above data comes from a normal population with a mean of 68 and standard deviation of 3.

[8 marks]

Sets, Relations, and Groups

7. [Maximum mark: 30]

(i) Let X and Y be two non-empty sets.

(a) Define the operation $X \bullet Y$ by $X \bullet Y = (X \cap Y) \cup (X' \cap Y')$. Prove that $(X \bullet Y)' = (X \cup Y) \cap (X' \cup Y')$. [3 marks]

(b) Let $f: \mathbb{N} \rightarrow \mathbb{N}$ be defined by $f(n) = n + 1$, for all $n \in \mathbb{N}$. Determine if f is an injection, a surjection, or a bijection. Give reasons for your answer. [3 marks]

(c) Let $h: X \rightarrow Y$, and let R be an equivalence relation on Y . $y_1 R y_2$ denotes that two elements y_1 and y_2 of Y are related.

Define a relation S on X by the following:

For all $a, b \in X$, $a S b$ if and only if $h(a) R h(b)$.

Determine if S is an equivalence relation on X . [4 marks]

(ii) (a) Let f_1, f_2, f_3, f_4 be functions defined on $\mathbb{Q} - \{0\}$, the set of rational numbers excluding zero, such that $f_1(z) = z$, $f_2(z) = -z$, $f_3(z) = \frac{1}{z}$, and

$$f_4(z) = -\frac{1}{z}, \text{ where } z \in \mathbb{Q} - \{0\}.$$

Let $T = \{f_1, f_2, f_3, f_4\}$. Define \circ as the composition of functions i.e. $(f_1 \circ f_2)(z) = f_1(f_2(z))$. Prove that (T, \circ) is an Abelian group. [6 marks]

(b) Let $G = \{1, 3, 5, 7\}$ and (G, \diamond) be the multiplicative group under the binary operation \diamond , multiplication modulo 8. Prove that the two groups (T, \circ) and (G, \diamond) are isomorphic. [5 marks]

(iii) Let a, b and p be elements of a group $(H, *)$ with an identity element e .

(a) If element a has order n and element a^{-1} has order m , then prove that $m = n$. [5 marks]

(b) If $b = p^{-1} * a * p$, prove, by mathematical induction, that $b^m = p^{-1} * a^m * p$, where $m = 1, 2, \dots$. [4 marks]

Discrete Mathematics


8. [Maximum marks: 30]

(i) A chemical manufacturer has to transport six chemical products from a factory to a processing plant by rail. Some of the products cannot be taken in the same railroad car because of the possibility of their mixing together and creating a violent reaction if an accident happens. The six products are labelled as A , B , C , D , E , and F . It is known that A cannot be kept in the same railroad car as B , C , or D ; B cannot be kept in the same railroad car as C or E ; C cannot be kept with D ; and E cannot be kept with F .

(a) Draw a graph where each vertex represents a product and the edges join pairs of products which cannot be in the same railroad car. [2 marks]

(b) Find the chromatic number of this graph and show a possible arrangement for transporting the six products. [4 marks]

(ii) Let G be the graph given below:



(a) Has G got an Eulerian circuit? Give a reason for your answer. [2 marks]

(b) What is the adjacency matrix of the graph G ? Determine how many walks of length 2 are there from vertex A to vertex C . [4 marks]

(c) Use Kruskal's algorithm to find the minimum spanning tree for graph G . [5 marks]

(iii) Find the solution of the difference equation

$$y_{n+2} = y_{n+1} + y_n, y_0 = 4, \text{ and } y_1 = 3. \quad [6 \text{ marks}]$$

(iv) (a) State the well-ordering principle for the set of positive integers. [2 marks]

(b) Extend this notion to any set where elements can be ordered. [2 marks]

(c) By giving reasons, carefully, establish whether \mathbb{Z} , the set of integers, is well-ordered. [3 marks]

Analysis and Approximations

9. [Maximum mark: 30]

(i) Let $f(x) = x^7 + 5x + 1$, $-2 \leq x \leq 2$.

(a) Use the Newton-Raphson method with an initial value $x_0 = -0.5$ to obtain a real root (zero) of f correct to 8 decimal places. Explain how your result for the third iteration has been obtained. [4 marks]

(b) Apply fixed point iteration with $x_0 = -0.5$ to calculate three iterates to find a solution of $f(x) = 0$. From your calculation, determine if the fixed point iteration will give a real root (zero) of f . Give a reason for your answer. [4 marks]

(c) Using Rolle's theorem prove that the equation $f(x) = 0$ has exactly one real root. [4 marks]

(ii) (a) Find the Maclaurin series of the function $g(x) = \sin x^2$ using the series expansion of $\sin x$, i.e. $\sin x = \sum_{n=0}^{\infty} (-1)^n \frac{x^{2n+1}}{(2n+1)!}$. [1 mark]

(b) Using the Maclaurin series of $g(x) = \sin x^2$ evaluate the definite integral

$$\int_0^1 \sin x^2 \, dx$$

correct to four decimal places. [5 marks]

(iii) (a) Use the ratio test to calculate the radius of convergence of the power series $\sum_{n=1}^{\infty} \frac{(x-5)^n}{n^{\frac{3}{2}}}$. [3 marks]

(b) Using your result from part (a), determine **all points** x where the power series given in (a) converges. [5 marks]

(iv) Using the mean value theorem prove that

$$|\sin x \cos x - \sin y \cos y| \leq |x - y|.$$
[4 marks]

Euclidean Geometry and Conic Sections

10. [Maximum mark: 30]

- (i) Let e_1 and e_2 be the eccentricities of the hyperbolas $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$ and $\frac{y^2}{b^2} - \frac{x^2}{a^2} = 1$, respectively. Prove that

$$\frac{1}{e_1^2} + \frac{1}{e_2^2} = 1 \quad [3 \text{ marks}]$$

- (ii) Let F_1 and F_2 be the foci of the hyperbola $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$ and P be any point on the hyperbola with coordinates (x_0, y_0) .

(a) Let (PM) be the tangent to the hyperbola at P , with M being on the x -axis between F_1 and F_2 . Find the coordinates of the point M . [4 marks]

(b) Find the lengths PF_1, PF_2, MF_1, MF_2 in terms of a, x_0 and e . [5 marks]

(c) Using an appropriate theorem from euclidean geometry, prove that $[PM]$ is the angle bisector of $F_1\hat{P}F_2$, stating which theorem you are using. [4 marks]

- (iii) If $ABCD$ is a cyclic quadrilateral, then prove Ptolemy's theorem:

$$AB \times CD + BC \times DA = AC \times BD \quad [5 \text{ marks}]$$

- (iv) In a triangle ABC , $\widehat{ACB} > \widehat{ABC}$. Let M and N be points on $[AC]$ and $[AB]$, respectively, such that $[CN]$ and $[BM]$ are the angle bisectors of \widehat{ACB} and \widehat{ABC} respectively. Let M' be the point on $[BM]$ which is nearer to the point M and is such that $M'\widehat{C}N = \frac{1}{2}(\widehat{ABC})$.

(a) Prove that $BNM'C$ is a cyclic quadrilateral. [4 marks]

(b) Prove that $BM' > CN$. [5 marks]

