



MARKSCHEME

November 2012

**MATHEMATICS
DISCRETE MATHEMATICS**

Higher Level

Paper 3

*This markscheme is **confidential** and for the exclusive use of examiners in this examination session.*

*It is the property of the International Baccalaureate and must **not** be reproduced or distributed to any other person without the authorization of the IB Assessment Centre.*

Instructions to Examiners

Abbreviations

- M** Marks awarded for attempting to use a correct **Method**; working must be seen.
- (M)** Marks awarded for **Method**; may be implied by **correct** subsequent working.
- A** Marks awarded for an **Answer** or for **Accuracy**; often dependent on preceding **M** marks.
- (A)** Marks awarded for an **Answer** or for **Accuracy**; may be implied by **correct** subsequent working.
- R** Marks awarded for clear **Reasoning**.
- N** Marks awarded for **correct** answers if **no** working shown.
- AG** Answer given in the question and so no marks are awarded.

Using the markscheme

1 General

Mark according to scoris instructions and the document “**Mathematics HL: Guidance for e-marking November 2012**”. It is **essential** that you read this document before you start marking. In particular, please note the following.

Marks must be recorded using the annotation stamps. Please check that you are entering marks for the right question.

- If a part is **completely correct**, (and gains all the “must be seen” marks), use the ticks with numbers to stamp full marks.
- If a part is completely wrong, stamp **A0** by the final answer.
- If a part gains anything else, it **must** be recorded using **all** the annotations.

All the marks will be added and recorded by scoris.

2 Method and Answer/Accuracy marks

- Do **not** automatically award full marks for a correct answer; all working **must** be checked, and marks awarded according to the markscheme.
- It is not possible to award **M0** followed by **AI**, as **A** mark(s) depend on the preceding **M** mark(s), if any.
- Where **M** and **A** marks are noted on the same line, *e.g.* **MIAI**, this usually means **MI** for an **attempt** to use an appropriate method (*e.g.* substitution into a formula) and **AI** for using the **correct** values.
- Where the markscheme specifies **(M2)**, **N3**, *etc.*, do **not** split the marks.
- Once a correct answer to a question or part-question is seen, ignore further working.

3 ***N* marks**

Award *N* marks for **correct** answers where there is **no** working.

- Do **not** award a mixture of *N* and other marks.
- There may be fewer *N* marks available than the total of *M*, *A* and *R* marks; this is deliberate as it penalizes candidates for not following the instruction to show their working.

4 **Implied marks**

Implied marks appear in **brackets e.g. (M1)**, and can only be awarded if **correct** work is seen or if implied in subsequent working.

- Normally the correct work is seen or implied in the next line.
- Marks **without** brackets can only be awarded for work that is **seen**.

5 **Follow through marks**

Follow through (**FT**) marks are awarded where an incorrect answer from one **part** of a question is used correctly in **subsequent** part(s). To award **FT** marks, **there must be working present** and not just a final answer based on an incorrect answer to a previous part.

- If the question becomes much simpler because of an error then use discretion to award fewer **FT** marks.
- If the error leads to an inappropriate value (e.g. $\sin \theta = 1.5$), do not award the mark(s) for the final answer(s).
- Within a question part, once an error is made, no further **dependent A** marks can be awarded, but **M** marks may be awarded if appropriate.
- Exceptions to this rule will be explicitly noted on the markscheme.

6 **Mis-read**

If a candidate incorrectly copies information from the question, this is a mis-read (**MR**). A candidate should be penalized only once for a particular mis-read. Use the **MR** stamp to indicate that this has been a mis-read. Then deduct the first of the marks to be awarded, even if this is an **M** mark, but award all others so that the candidate only loses one mark.

- If the question becomes much simpler because of the **MR**, then use discretion to award fewer marks.
- If the **MR** leads to an inappropriate value (e.g. $\sin \theta = 1.5$), do not award the mark(s) for the final answer(s).

7 **Discretionary marks (d)**

An examiner uses discretion to award a mark on the rare occasions when the markscheme does not cover the work seen. In such cases the annotation **DM** should be used and a brief **note** written next to the mark explaining this decision.

8 Alternative methods

Candidates will sometimes use methods other than those in the markscheme. Unless the question specifies a method, other correct methods should be marked in line with the markscheme. If in doubt, contact your team leader for advice.

- Alternative methods for complete questions are indicated by **METHOD 1**, **METHOD 2**, etc.
- Alternative solutions for part-questions are indicated by **EITHER . . . OR**.
- Where possible, alignment will also be used to assist examiners in identifying where these alternatives start and finish.

9 Alternative forms

Unless the question specifies otherwise, **accept** equivalent forms.

- As this is an international examination, accept all alternative forms of **notation**.
- In the markscheme, equivalent **numerical** and **algebraic** forms will generally be written in brackets immediately following the answer.
- In the markscheme, **simplified** answers, (which candidates often do not write in examinations), will generally appear in brackets. Marks should be awarded for either the form preceding the bracket or the form in brackets (if it is seen).

Example: for differentiating $f(x) = 2 \sin(5x - 3)$, the markscheme gives:

$$f'(x) = (2 \cos(5x - 3)) 5 \quad (= 10 \cos(5x - 3)) \quad \text{AI}$$

Award **AI** for $(2 \cos(5x - 3)) 5$, even if $10 \cos(5x - 3)$ is not seen.

10 Accuracy of Answers

Candidates should **NO LONGER** be penalized for an accuracy error (**AP**).

If the level of accuracy is specified in the question, a mark will be allocated for giving the answer to the required accuracy. When this is not specified in the question, all numerical answers should be given exactly or correct to three significant figures. Please check work carefully for **FT**.

11 Crossed out work

If a candidate has drawn a line through work on their examination script, or in some other way crossed out their work, do not award any marks for that work.

12 Calculators

A GDC is required for paper 3, but calculators with symbolic manipulation features (e.g. TI-89) are not allowed.

Calculator notation

The Mathematics HL guide says:


Students must always use correct mathematical notation, not calculator notation.

Do **not** accept final answers written using calculator notation. However, do not penalize the use of calculator notation in the working.

13 More than one solution

Where a candidate offers two or more different answers to the same question, an examiner should only mark the first response unless the candidate indicates otherwise.

1. (a)



from tabular method as shown above (or equivalent)

M1A1A1

Note: Award the first *AI* for obtaining 3 as the shortest distance to C.
Award the second *AI* for obtaining the rest of the shortest distances.

shortest path has length 17

AI

backtracking as shown above (or equivalent)

(M1)

shortest path is SABT

AI

[6 marks]

(b) (i) no, as *S* and *T* have odd degree

A1R1

Note: Mentioning one vertex of odd degree is sufficient.

(ii) yes, as only *S* and *T* have odd degree

A1R1

Note: In each case only award the *AI* if the *R1* has been given.
Accept an actual trail in (b)(ii).

[4 marks]

(c) Huan has to travel along all the edges via an open Eulerian trail of length
 $4+3+5+2+1+3+5+4+7+8+5+6+7+6+6+8+9=94$
 and then back to *S* from *T* along the shortest path found in (a) of length 17
 so shortest total distance is $94+17=111$

R1

AI

R1

AI

[4 marks]

Total [14 marks]

2. (a)




AIAIAIAI

Note: Award *AI* for 1 correct graph, then *AI* for each of the subsequent 3 graphs, provided they are not repeated.

[4 marks]

(b)



AIAIAIAIAIAI

Note: Award *AI* for 1 correct tree, then *AI* for each of the subsequent 5 trees, provided they are not repeated.

[6 marks]

Total [10 marks]

3. (a) **METHOD 1**

861	957	[1, 0]	[0, 1]
	-1×861	$\frac{-861}{96}$	$\frac{-[1, 0]}{[-1, 1]}$
-8×96	$\frac{-768}{93}$	$\frac{-8[-1, 1]}{[9, -8]}$	
	-1×93	$\frac{-93}{3}$	$\frac{-[9, -8]}{[-10, 9]}$

by the above working on the left (or similar)

MIAIAI

Note: Award *AI* for 96 and *AI* for 93.

$h = 3$ (since 3 divides 93)

AI

[4 marks]

METHOD 2

$957 = 861 + 96$

MIAI

$861 = 8 \times 96 + 93$

AI

$96 = 93 + 3$

so $h = 3$ (since 3 divides 93)

AI

[4 marks]

(b) **METHOD 1**

if method 1 was used for part (a)

by the above working on the right (or equivalent)

MI

$-10 \times 861 + 9 \times 957 = 3$

so $A = -10$ and $B = 9$

AIAI

[3 marks]

METHOD 2

$3 = 96 - 93$

$= 96 - (861 - 8 \times 96) = 9 \times 96 - 861$

MI

$= 9 \times (957 - 861) - 861$

$= -10 \times 861 + 9 \times 957$

so $A = -10$ and $B = 9$

AIAI

[3 marks]

(c) from (b) $-10 \times 287 + 9 \times 319 = 1$ so

AI

$-10 \times 287 \equiv 1 \pmod{319}$

AI

$287w \equiv 2 \pmod{319} \Rightarrow -10 \times 287w \equiv -10 \times 2 \pmod{319}$

MI

$\Rightarrow w \equiv -20 \pmod{319}$

AI

so $w = 299$

AI

[5 marks]

continued ...

Question 3 continued

- (d) from (b) $-10 \times 861 + 9 \times 957 = 3 \Rightarrow -20 \times 861 + 18 \times 957 = 6$
 so general solution is $x = -20 + 319t$
 $y = 18 - 287t \ (t \in \mathbb{Z})$

M1A1
A1A1
A1A1

[6 marks]

Total [18 marks]

- 4. (a) if p is a prime (and $a \not\equiv 0 \pmod{p}$) with $a \in \mathbb{Z}$) then
 $a^{p-1} \equiv 1 \pmod{p}$

A1
A1

[2 marks]

Note: Accept $a^p \equiv a \pmod{p}$.

- (b) by Fermat $x^4 \equiv 1 \pmod{5}$
 so $x^{12} \equiv 1 \pmod{5}$
 by Fermat $x^{12} \equiv 1 \pmod{13}$
 so $x^{12} \equiv 1 \pmod{65}$
 let $x^{12} = 1 + 65t$, then $1 + 65t \equiv 4 \pmod{11}$
 so $-t \equiv 3 \pmod{11} \Rightarrow t \equiv -3 \pmod{11}$
 $\Rightarrow t \equiv 8 \pmod{11}$
 $x^{12} = 1 + 65(8 + 11w) = 521 + 715w$
 so remainder is 521

A1
A1
A1
A1
M1

A1

M1

A1


Note: For solutions that start by showing that x^2 is congruent to 4 mod 11 award suitable discretion marks being careful that the solutions are logical and cover all possibilities.
--

[8 marks]


Total [10 marks]

5. G must have at least 3 edges *RI*
 since G is connected and so it must have a spanning tree as a subgraph and also $V = 4$
 similarly \overline{G} must have at least 3 edges *RI*
 but K_4 has 6 edges so both G and \overline{G} have exactly 3 edges *RI*
 there are two possibilities for G


case (i)



case (ii)




in case (ii) \overline{G} is




this is not connected so case (ii) cannot happen *RI*

in case (i) \overline{G} is



this is isomorphic to



so G and \overline{G} are isomorphic *AI*

AI

AG

Note: There are other equally valid proofs for example considering the possible degrees of the vertices.

[8 marks]