

Markscheme

November 2017

Mathematics

Higher level

Paper 1

This markscheme is the property of the International Baccalaureate and must **not** be reproduced or distributed to any other person without the authorization of the IB Global Centre, Cardiff.

Instructions to Examiners

Abbreviations

- M** Marks awarded for attempting to use a valid **Method**; working must be seen.
- (M)** Marks awarded for **Method**; may be implied by **correct** subsequent working.
- A** Marks awarded for an **Answer** or for **Accuracy**; often dependent on preceding **M** marks.
- (A)** Marks awarded for an **Answer** or for **Accuracy**; may be implied by **correct** subsequent working.
- R** Marks awarded for clear **Reasoning**.
- N** Marks awarded for **correct** answers if **no** working shown.
- AG** Answer given in the question and so no marks are awarded.

Using the markscheme

1 General

Mark according to RM™ Assessor instructions and the document “**Mathematics HL: Guidance for e-marking November 2017**”. It is essential that you read this document before you start marking.

In particular, please note the following:

- Marks must be recorded using the annotation stamps. Please check that you are entering marks for the right question.
- If a part is **completely correct**, (and gains all the “must be seen” marks), use the ticks with numbers to stamp full marks.
- If a part is completely wrong, stamp **A0** by the final answer.
- If a part gains anything else, it **must** be recorded using **all** the annotations.
- All the marks will be added and recorded by RM™ Assessor.

2 Method and Answer/Accuracy marks

- Do **not** automatically award full marks for a correct answer; all working **must** be checked, and marks awarded according to the markscheme.
- It is not possible to award **M0** followed by **A1**, as **A** mark(s) depend on the preceding **M** mark(s), if any.
- Where **M** and **A** marks are noted on the same line, eg **M1A1**, this usually means **M1** for an **attempt** to use an appropriate method (eg substitution into a formula) and **A1** for using the **correct** values.
- Where the markscheme specifies (**M2**), **N3**, etc., do **not** split the marks.

- Once a correct answer to a question or part-question is seen, ignore further correct working. However, if further working indicates a lack of mathematical understanding do not award the final **A1**. An exception to this may be in numerical answers, where a correct exact value is followed by an incorrect decimal. However, if the incorrect decimal is carried through to a subsequent part, and correct **FT** working shown, award **FT** marks as appropriate but do not award the final **A1** in that part.

Examples

	Correct answer seen	Further working seen	Action
1.	$8\sqrt{2}$	5.65685... (incorrect decimal value)	Award the final A1 (ignore the further working)
2.	$\frac{1}{4}\sin 4x$	$\sin x$	Do not award the final A1
3.	$\log a - \log b$	$\log(a - b)$	Do not award the final A1

3 **N** marks

Award **N** marks for **correct** answers where there is **no** working.

- Do **not** award a mixture of **N** and other marks.
- There may be fewer **N** marks available than the total of **M**, **A** and **R** marks; this is deliberate as it penalizes candidates for not following the instruction to show their working.

4 **Implied marks**

*Implied marks appear in **brackets eg (M1)**, and can only be awarded if **correct** work is seen or if implied in subsequent working.*

- Normally the correct work is seen or implied in the next line.
- Marks **without** brackets can only be awarded for work that is **seen**.

5 **Follow through marks**

*Follow through (**FT**) marks are awarded where an incorrect answer from one **part** of a question is used correctly in **subsequent** part(s). To award **FT** marks, **there must be working present** and not just a final answer based on an incorrect answer to a previous part.*

- If the question becomes much simpler because of an error then use discretion to award fewer **FT** marks.
- If the error leads to an inappropriate value (eg $\sin \theta = 1.5$), do not award the mark(s) for the final answer(s).
- Within a question part, once an error is made, no further **dependent A** marks can be awarded, but **M** marks may be awarded if appropriate.
- Exceptions to this rule will be explicitly noted on the markscheme.

6 Misread

If a candidate incorrectly copies information from the question, this is a misread (**MR**). A candidate should be penalized only once for a particular misread. Use the **MR** stamp to indicate that this has been a misread. Then deduct the first of the marks to be awarded, even if this is an **M** mark, but award all others so that the candidate only loses **[1 mark]**.

- If the question becomes much simpler because of the **MR**, then use discretion to award fewer marks.
- If the **MR** leads to an inappropriate value (eg $\sin \theta = 1.5$), do not award the mark(s) for the final answer(s).

7 Discretionary marks (**d**)

An examiner uses discretion to award a mark on the rare occasions when the markscheme does not cover the work seen. In such cases the annotation **DM** should be used and a brief **note** written next to the mark explaining this decision.

8 Alternative methods

Candidates will sometimes use methods other than those in the markscheme. Unless the question specifies a method, other correct methods should be marked in line with the markscheme. If in doubt, contact your team leader for advice.

- Alternative methods for complete questions are indicated by **METHOD 1**, **METHOD 2**, etc.
- Alternative solutions for part-questions are indicated by **EITHER . . . OR**.
- Where possible, alignment will also be used to assist examiners in identifying where these alternatives start and finish.

9 Alternative forms

Unless the question specifies otherwise, **accept** equivalent forms.

- As this is an international examination, accept all alternative forms of **notation**.
- In the markscheme, equivalent **numerical** and **algebraic** forms will generally be written in brackets immediately following the answer.
- In the markscheme, **simplified** answers, (which candidates often do not write in examinations), will generally appear in brackets. Marks should be awarded for either the form preceding the bracket or the form in brackets (if it is seen).

Example: for differentiating $f(x) = 2\sin(5x - 3)$, the markscheme gives

$$f'(x) = (2\cos(5x - 3))5 \quad (=10\cos(5x - 3)) \quad \mathbf{A1}$$

Award **A1** for $(2\cos(5x - 3))5$, even if $10\cos(5x - 3)$ is not seen.

10 Accuracy of Answers

Candidates should **NO LONGER** be penalized for an accuracy error (**AP**).

*If the level of accuracy is specified in the question, a mark will be allocated for giving the answer to the required accuracy. When this is not specified in the question, all numerical answers should be given exactly or correct to three significant figures. Please check work carefully for **FT**.*

11 Crossed out work

If a candidate has drawn a line through work on their examination script, or in some other way crossed out their work, do not award any marks for that work.

12 Calculators

No calculator is allowed. The use of any calculator on paper 1 is malpractice, and will result in no grade awarded. If you see work that suggests a candidate has used any calculator, please follow the procedures for malpractice. Examples: finding an angle, given a trig ratio of 0.4235.

13 More than one solution

Where a candidate offers two or more different answers to the same question, an examiner should only mark the first response unless the candidate indicates otherwise.

14. Candidate work

Candidates are meant to write their answers to Section A on the question paper (QP), and Section B on answer booklets. Sometimes, they need more room for Section A, and use the booklet (and often comment to this effect on the QP), or write outside the box. This work should be marked.

The instructions tell candidates not to write on Section B of the QP. Thus they may well have done some rough work here which they assume will be ignored. If they have solutions on the answer booklets, there is no need to look at the QP. However, if there are whole questions or whole part solutions missing on answer booklets, please check to make sure that they are not on the QP, and if they are, mark those whole questions or whole part solutions that have not been written on answer booklets.

Section A

1. $\log_2(x + 3) + \log_2(x - 3) = 4$

$\log_2(x^2 - 9) = 4$ (M1)

$x^2 - 9 = 2^4 (=16)$ M1A1

$x^2 = 25$

$x = \pm 5$ (A1)

$x = 5$ A1

[5 marks]

2. (a) $\vec{AB} = \begin{pmatrix} 6 \\ -8 \\ 17 \end{pmatrix}$ (A1)

$r = \begin{pmatrix} 0 \\ 3 \\ -6 \end{pmatrix} + \lambda \begin{pmatrix} 6 \\ -8 \\ 17 \end{pmatrix}$ or $r = \begin{pmatrix} 6 \\ -5 \\ 11 \end{pmatrix} + \lambda \begin{pmatrix} 6 \\ -8 \\ 17 \end{pmatrix}$ M1A1

Note: Award **M1A0** if $r =$ is not seen (or equivalent).

[3 marks]

(b) substitute line L in $\Pi : 4(6\lambda) - 3(3 - 8\lambda) + 2(-6 + 17\lambda) = 20$ M1

$82\lambda = 41$

$\lambda = \frac{1}{2}$ (A1)

$r = \begin{pmatrix} 0 \\ 3 \\ -6 \end{pmatrix} + \frac{1}{2} \begin{pmatrix} 6 \\ -8 \\ 17 \end{pmatrix} = \begin{pmatrix} 3 \\ -1 \\ \frac{5}{2} \end{pmatrix}$

so coordinate is $\left(3, -1, \frac{5}{2}\right)$ A1

Note: Accept coordinate expressed as position vector

$$\begin{pmatrix} 3 \\ -1 \\ \frac{5}{2} \end{pmatrix} .$$

[3 marks]

Total [6 marks]

3. (a) $q(4) = 0$ (M1)
 $192 - 176 + 4k + 8 = 0$ ($24 + 4k = 0$) A1
 $k = -6$ A1
[3 marks]

- (b) $3x^3 - 11x^2 - 6x + 8 = (x - 4)(3x^2 + px - 2)$
 equate coefficients of x^2 : (M1)
 $-12 + p = -11$
 $p = 1$
 $(x - 4)(3x^2 + x - 2)$ (A1)
 $(x - 4)(3x - 2)(x + 1)$ A1

Note: Allow part (b) marks if any of this work is seen in part (a).

Note: Allow equivalent methods (eg, synthetic division) for the **M** marks in each part.

[3 marks]

Total [6 marks]

4. each term is of the form $\binom{7}{r}(x^2)^{7-r}\left(\frac{-2}{x}\right)^r$ (M1)
 $= \binom{7}{r}x^{14-2r}(-2)^r x^{-r}$
 so $14 - 3r = 8$ (A1)
 $r = 2$
 so require $\binom{7}{2}(x^2)^5\left(\frac{-2}{x}\right)^2$ (or simply $\binom{7}{2}(-2)^2$) A1
 $= 21 \times 4$
 $= 84$ A1

Note: Candidates who attempt a full expansion, including the correct term, may only be awarded **M1A0A0A0**.

[4 marks]

5. $s = \int_0^{\frac{1}{2}} 10te^{-2t} dt$

attempt at integration by parts

M1

$$= \left[-5te^{-2t} \right]_0^{\frac{1}{2}} - \int_0^{\frac{1}{2}} -5e^{-2t} dt$$

A1

$$= \left[-5te^{-2t} - \frac{5}{2}e^{-2t} \right]_0^{\frac{1}{2}}$$

(A1)

Note: Condone absence of limits (or incorrect limits) and missing factor of 10 up to this point.

$$s = \int_0^{\frac{1}{2}} 10te^{-2t} dt$$


(M1)

$$= -5e^{-1} + \frac{5}{2} \left(= \frac{-5}{e} + \frac{5}{2} \right) \left(= \frac{5e-10}{2e} \right)$$

A1

[5 marks]

6. (a)



correct vertical asymptote
shape including correct horizontal asymptote

$$\left(\frac{1}{3}, 0\right)$$

$$\left(0, -\frac{1}{2}\right)$$

A1

A1

A1

A1


Note: Accept $x = \frac{1}{3}$ and $y = -\frac{1}{2}$ marked on the axes.

[4 marks]

continued...

Question 6 continued

(b) **METHOD 1**



$$\frac{1 - 3x}{x - 2} = 2$$

(M1)

$$\Rightarrow x = 1$$

A1

$$-\left(\frac{1 - 3x}{x - 2}\right) = 2$$

(M1)

Note: Award this **M1** for the line above or a correct sketch identifying a second critical value.

$$\Rightarrow x = -3$$

A1

$$\text{solution is } -3 < x < 1$$

A1

[5 marks]

METHOD 2

$$|1 - 3x| < 2|x - 2|, x \neq 2$$

$$1 - 6x + 9x^2 < 4(x^2 - 4x + 4)$$

(M1)A1

$$1 - 6x + 9x^2 < 4x^2 - 16x + 16$$

$$5x^2 + 10x - 15 < 0$$

$$x^2 + 2x - 3 < 0$$

A1

$$(x + 3)(x - 1) < 0$$

(M1)

$$\text{solution is } -3 < x < 1$$

A1

[5 marks]

continued...

Question 6 continued

METHOD 3

$$-2 < \frac{1-3x}{x-2} < 2$$

consider $\frac{1-3x}{x-2} < 2$ **(M1)**

Note: Also allow consideration of “>” or “=” for the awarding of the **M** mark.

recognition of critical value at $x = 1$ **A1**

consider $-2 < \frac{1-3x}{x-2}$ **(M1)**

Note: Also allow consideration of “>” or “=” for the awarding of the **M** mark.

recognition of critical value at $x = -3$ **A1**

solution is $-3 < x < 1$ **A1**

[5 marks]

Total [9 marks]

7. $x^3 + y^3 - 3xy = 0$

$$3x^2 + 3y^2 \frac{dy}{dx} - 3x \frac{dy}{dx} - 3y = 0$$
 M1A1

Note: Differentiation wrt y is also acceptable.

$$\frac{dy}{dx} = \frac{3y - 3x^2}{3y^2 - 3x} \left(= \frac{y - x^2}{y^2 - x} \right)$$
 (A1)

Note: All following marks may be awarded if the denominator is correct, but the numerator incorrect.

$$y^2 - x = 0$$
 M1

EITHER

$$x = y^2$$

$$y^6 + y^3 - 3y^3 = 0$$
 M1A1

$$y^6 - 2y^3 = 0$$

$$y^3(y^3 - 2) = 0$$

$$(y \neq 0) \therefore y = \sqrt[3]{2}$$
 A1

$$x = (\sqrt[3]{2})^2 (= \sqrt[3]{4})$$
 A1

continued...

Question 7 continued

OR

$$x^3 + xy - 3xy = 0$$

M1

$$x(x^2 - 2y) = 0$$

$$x \neq 0 \Rightarrow y = \frac{x^2}{2}$$

A1

$$y^2 = \frac{x^4}{4}$$

$$x = \frac{x^4}{4}$$

$$x(x^3 - 4) = 0$$

$$(x \neq 0) \therefore x = \sqrt[3]{4}$$

A1

$$y = \frac{(\sqrt[3]{4})^2}{2} = \sqrt[3]{2}$$

A1

[8 marks]

8. METHOD 1

$$216i = 216 \left(\cos \frac{\pi}{2} + i \sin \frac{\pi}{2} \right)$$

A1

$$z + 2i = \sqrt[3]{216} \left(\cos \left(\frac{\pi}{2} + 2\pi k \right) + i \sin \left(\frac{\pi}{2} + 2\pi k \right) \right)^{\frac{1}{3}}$$

(M1)

$$z + 2i = 6 \left(\cos \left(\frac{\pi}{6} + \frac{2\pi k}{3} \right) + i \sin \left(\frac{\pi}{6} + \frac{2\pi k}{3} \right) \right)$$

A1

$$z_1 + 2i = 6 \left(\cos \frac{\pi}{6} + i \sin \frac{\pi}{6} \right) = 6 \left(\frac{\sqrt{3}}{2} + \frac{i}{2} \right) = 3\sqrt{3} + 3i$$

$$z_2 + 2i = 6 \left(\cos \frac{5\pi}{6} + i \sin \frac{5\pi}{6} \right) = 6 \left(\frac{-\sqrt{3}}{2} + \frac{i}{2} \right) = -3\sqrt{3} + 3i$$

$$z_3 + 2i = 6 \left(\cos \frac{3\pi}{2} + i \sin \frac{3\pi}{2} \right) = -6i$$

A2

Note: Award **A1A0** for one correct root.

so roots are $z_1 = 3\sqrt{3} + i$, $z_2 = -3\sqrt{3} + i$ and $z_3 = -8i$

M1A1

Note: Award **M1** for subtracting $2i$ from their three roots.

[7 marks]

continued...

Question 8 continued

METHOD 2

$$(a\sqrt{3} + (b + 2)i)^3 = 216i$$

$$(a\sqrt{3})^3 + 3(a\sqrt{3})^2(b + 2)i - 3(a\sqrt{3})(b + 2)^2 - i(b + 2)^3 = 216i \quad \mathbf{M1A1}$$

$$(a\sqrt{3})^3 - 3(a\sqrt{3})(b + 2)^2 + i(3(a\sqrt{3})^2(b + 2) - (b + 2)^3) = 216i$$

$$(a\sqrt{3})^3 - 3(a\sqrt{3})(b + 2)^2 = 0 \text{ and } 3(a\sqrt{3})^2(b + 2) - (b + 2)^3 = 216 \quad \mathbf{M1A1}$$

$$a(a^2 - (b + 2)^2) = 0 \text{ and } 9a^2(b + 2) - (b + 2)^3 = 216$$

$$a = 0 \text{ or } a^2 = (b + 2)^2$$

$$\text{if } a = 0, -(b + 2)^3 = 216 \Rightarrow b + 2 = -6$$

$$\therefore b = -8 \quad \mathbf{A1}$$

$$(a, b) = (0, -8)$$

$$\text{if } a^2 = (b + 2)^2, 9(b + 2)^2(b + 2) - (b + 2)^3 = 216$$

$$8(b + 2)^3 = 216$$

$$(b + 2)^3 = 27$$

$$b + 2 = 3$$

$$b = 1$$

$$\therefore a^2 = 9 \Rightarrow a = \pm 3$$

$$\therefore (a, b) = (\pm 3, 1) \quad \mathbf{A1A1}$$

$$\text{so roots are } z_1 = 3\sqrt{3} + i, z_2 = -3\sqrt{3} + i \text{ and } z_3 = -8i$$

continued...

Question 8 continued

METHOD 3

$$(z + 2i)^3 - (-6i)^3 = 0$$

attempt to factorise:

$$((z + 2i) - (-6i))((z + 2i)^2 + (z + 2i)(-6i) + (-6i)^2) = 0$$

$$(z + 8i)(z^2 - 2iz - 28) = 0$$

$$z + 8i = 0 \Rightarrow z = -8i$$

$$z^2 - 2iz - 28 = 0 \Rightarrow z = \frac{2i \pm \sqrt{-4 - (4 \times 1 \times -28)}}{2}$$

$$z = \frac{2i \pm \sqrt{108}}{2}$$

$$z = \frac{2i \pm 6\sqrt{3}}{2}$$

$$z = i \pm 3\sqrt{3}$$

M1

A1

A1

A1

M1

A1A1

Special Case:

Note: If a candidate recognises that $\sqrt[3]{216i} = -6i$ (anywhere seen), and makes no valid progress in finding three roots, award **A1** only.

[7 marks]

Section B

9. (a) (i) $\vec{OF} = \frac{1}{7}\mathbf{b}$ **A1**
- (ii) $\vec{AF} = \vec{OF} - \vec{OA}$ **(M1)**
 $= \frac{1}{7}\mathbf{b} - \mathbf{a}$ **A1**
- [3 marks]**
- (b) (i) $\vec{OD} = \mathbf{a} + \lambda\left(\frac{1}{7}\mathbf{b} - \mathbf{a}\right) \left(= (1 - \lambda)\mathbf{a} + \frac{\lambda}{7}\mathbf{b}\right)$ **M1A1**
- (ii) $\vec{OD} = \frac{1}{2}\mathbf{a} + \mu\left(-\frac{1}{2}\mathbf{a} + \mathbf{b}\right) \left(= \left(\frac{1}{2} - \frac{\mu}{2}\right)\mathbf{a} + \mu\mathbf{b}\right)$ **M1A1**
- [4 marks]**
- (c) equating coefficients: **M1**
 $\frac{\lambda}{7} = \mu, 1 - \lambda = \frac{1 - \mu}{2}$ **A1**
solving simultaneously: **M1**
 $\lambda = \frac{7}{13}, \mu = \frac{1}{13}$ **A1AG**
- [4 marks]**
- (d) $\vec{CD} = \frac{1}{13}\vec{CB}$
 $= \frac{1}{13}\left(\mathbf{b} - \frac{1}{2}\mathbf{a}\right) \left(= -\frac{1}{26}\mathbf{a} + \frac{1}{13}\mathbf{b}\right)$ **M1A1**
- [2 marks]**

continued...

Question 9 continued

(e) **METHOD 1**

$$\text{area } \triangle ACD = \frac{1}{2} CD \times AC \times \sin \hat{A}CB \quad (M1)$$

$$\text{area } \triangle ACB = \frac{1}{2} CB \times AC \times \sin \hat{A}CB \quad (M1)$$

$$\text{ratio } \frac{\text{area } \triangle ACD}{\text{area } \triangle ACB} = \frac{CD}{CB} = \frac{1}{13} \quad A1$$

$$k = \frac{\text{area } \triangle OAB}{\text{area } \triangle CAD} = \frac{13}{\text{area } \triangle CAB} \times \text{area } \triangle OAB \quad (M1)$$

$$= 13 \times 2 = 26 \quad A1$$

METHOD 2

$$\text{area } \triangle OAB = \frac{1}{2} |\mathbf{a} \times \mathbf{b}| \quad A1$$

$$\text{area } \triangle CAD = \frac{1}{2} \left| \vec{CA} \times \vec{CD} \right| \text{ or } \frac{1}{2} \left| \vec{CA} \times \vec{AD} \right| \quad M1$$

$$= \frac{1}{2} \left| \frac{1}{2} \mathbf{a} \times \left(-\frac{1}{26} \mathbf{a} + \frac{1}{13} \mathbf{b} \right) \right|$$

$$= \frac{1}{2} \left| \frac{1}{2} \mathbf{a} \times \left(-\frac{1}{26} \mathbf{a} \right) + \frac{1}{2} \mathbf{a} \times \frac{1}{13} \mathbf{b} \right| \quad (M1)$$

$$= \frac{1}{2} \times \frac{1}{2} \times \frac{1}{13} |\mathbf{a} \times \mathbf{b}| \left(= \frac{1}{52} |\mathbf{a} \times \mathbf{b}| \right) \quad A1$$

$$\text{area } \triangle OAB = k(\text{area } \triangle CAD)$$

$$\frac{1}{2} |\mathbf{a} \times \mathbf{b}| = k \frac{1}{52} |\mathbf{a} \times \mathbf{b}|$$

$$k = 26 \quad A1$$

[5 marks]

Total [18 marks]

10. (a) **METHOD 1**

$$\text{number of possible "deals"} = 4! = 24 \quad A1$$

consider ways of achieving "no matches" (Chloe winning):

Selena could deal B, C, D (ie, 3 possibilities)

as her first card R1

for each of these matches, there are only 3 possible combinations for the remaining 3 cards R1

$$\text{so no. ways achieving no matches} = 3 \times 3 = 9 \quad M1A1$$

$$\text{so probability Chloe wins} = \frac{9}{24} = \frac{3}{8} \quad A1AG$$

continued...

Question 10 continued

METHOD 2

number of possible “deals” = $4! = 24$ **A1**

consider ways of achieving a match (Selena winning)

Selena card A can match with Chloe card A, giving 6 possibilities for this happening **R1**

if Selena deals B as her first card, there are only 3 possible combinations for the remaining 3 cards. Similarly for dealing C and dealing D **R1**

so no. ways achieving one match is $= 6 + 3 + 3 + 3 = 15$ **M1A1**

so probability Chloe wins $= 1 - \frac{15}{24} = \frac{3}{8}$ **A1AG**

METHOD 3

systematic attempt to find number of outcomes where Chloe wins (no matches) (using tree diag. or otherwise) **M1**

9 found **A1**

each has probability $\frac{1}{4} \times \frac{1}{3} \times \frac{1}{2} \times 1$ **M1**

$= \frac{1}{24}$ **A1**

their 9 multiplied by their $\frac{1}{24}$ **M1A1**

$= \frac{3}{8}$ **AG**

[6 marks]

continued...

Question 10 continued

(b) (i) $X \sim B\left(50, \frac{3}{8}\right)$ (M1)

$$\mu = np = 50 \times \frac{3}{8} = \frac{150}{8} \left(= \frac{75}{4} \right) (= 18.75)$$
 (M1)A1

(ii) $\sigma^2 = np(1-p) = 50 \times \frac{3}{8} \times \frac{5}{8} = \frac{750}{64} \left(= \frac{375}{32} \right) (= 11.7)$ (M1)A1

[5 marks]

Total [11 marks]

11. (a) even function A1
 since $\cos kx = \cos(-kx)$ **and** $f_n(x)$ is a product of even functions R1

OR

even function A1
 since $(\cos 2x)(\cos 4x)\dots = (\cos(-2x))(\cos(-4x))\dots$ R1

Note: Do not award **A0R1**.

[2 marks]

(b) consider the case $n = 1$

$$\frac{\sin 4x}{2 \sin 2x} = \frac{2 \sin 2x \cos 2x}{2 \sin 2x} = \cos 2x$$
 M1
 hence true for $n = 1$ R1

assume true for $n = k$, ie, $(\cos 2x)(\cos 4x)\dots(\cos 2^k x) = \frac{\sin 2^{k+1} x}{2^k \sin 2x}$ M1

Note: Do not award **M1** for “let $n = k$ ” or “assume $n = k$ ” or equivalent.

consider $n = k + 1$:

$$f_{k+1}(x) = f_k(x)(\cos 2^{k+1} x)$$
 (M1)

$$= \frac{\sin 2^{k+1} x}{2^k \sin 2x} \cos 2^{k+1} x$$
 A1

$$= \frac{2 \sin 2^{k+1} x \cos 2^{k+1} x}{2^{k+1} \sin 2x}$$
 A1

$$= \frac{\sin 2^{k+2} x}{2^{k+1} \sin 2x}$$
 A1

so $n = 1$ true and $n = k$ true $\Rightarrow n = k + 1$ true. Hence true for all $n \in \mathbb{Z}^+$ R1

Note: To obtain the final **R1**, all the previous **M** marks must have been awarded.

[8 marks]

continued...

Question 11 continued

(c) attempt to use $f' = \frac{vu' - uv'}{v^2}$ (or correct product rule) **M1**

$$f'_n(x) = \frac{(2^n \sin 2x)(2^{n+1} \cos 2^{n+1} x) - (\sin 2^{n+1} x)(2^{n+1} \cos 2x)}{(2^n \sin 2x)^2} \quad \text{A1A1}$$

Note: Award **A1** for correct numerator and **A1** for correct denominator.

[3 marks]

(d) $f'_n\left(\frac{\pi}{4}\right) = \frac{\left(2^n \sin \frac{\pi}{2}\right)\left(2^{n+1} \cos 2^{n+1} \frac{\pi}{4}\right) - \left(\sin 2^{n+1} \frac{\pi}{4}\right)\left(2^{n+1} \cos \frac{\pi}{2}\right)}{\left(2^n \sin \frac{\pi}{2}\right)^2} \quad \text{(M1)(A1)}$

$$f'_n\left(\frac{\pi}{4}\right) = \frac{(2^n)\left(2^{n+1} \cos 2^{n+1} \frac{\pi}{4}\right)}{(2^n)^2} \quad \text{(A1)}$$

$$= 2 \cos 2^{n+1} \frac{\pi}{4} (= 2 \cos 2^{n-1} \pi) \quad \text{A1}$$

$$f'_n\left(\frac{\pi}{4}\right) = 2 \quad \text{A1}$$

$$f_n\left(\frac{\pi}{4}\right) = 0 \quad \text{A1}$$

Note: This **A** mark is independent from the previous marks.

$$y = 2\left(x - \frac{\pi}{4}\right) \quad \text{M1A1}$$

$$4x - 2y - \pi = 0 \quad \text{AG}$$

[8 marks]

Total [21 marks]