

IB Maths DP

YOUR NOTES



4. Statistics & Probability

CONTENTS

- 4.1 Statistics Toolkit
 - 4.1.1 Sampling & Data Collection
 - 4.1.2 Statistical Measures
 - 4.1.3 Frequency Tables
 - 4.1.4 Linear Transformations of Data
 - 4.1.5 Outliers
 - 4.1.6 Univariate Data
 - 4.1.7 Interpreting Data
- 4.2 Correlation & Regression
 - 4.2.1 Bivariate data
 - 4.2.2 Correlation Coefficients
 - 4.2.3 Linear Regression
- 4.3 Probability
 - 4.3.1 Probability & Types of Events
 - 4.3.2 Conditional Probability
 - 4.3.3 Sample Space Diagrams
- 4.4 Probability Distributions
 - 4.4.1 Discrete Probability Distributions
 - 4.4.2 Expected Values
- 4.5 Binomial Distribution
 - 4.5.1 The Binomial Distribution
 - 4.5.2 Calculating Binomial Probabilities
- 4.6 Normal Distribution
 - 4.6.1 The Normal Distribution
 - 4.6.2 Calculations with Normal Distribution
- 4.7 Hypothesis Testing
 - 4.7.1 Hypothesis Testing
 - 4.7.2 Chi-squared Test for Independence
 - 4.7.3 Goodness of Fit Test
 - 4.7.4 The t-test

4.1 Statistics Toolkit

4.1.1 Sampling & Data Collection

YOUR NOTES



Types of Data

What are the different types of data?

- **Qualitative** data is data that is usually given in words not numbers to **describe** something
 - For example: the colour of a teacher's car
- **Quantitative** data is data that is given using numbers which **counts or measures** something
 - For example: the number of pets that a student has
- **Discrete** data is quantitative data that needs to be **counted**
 - Discrete data can only take **specific values** from a set of (usually finite) values
 - For example: the number of times a coin is flipped until a 'tails' is obtained
- **Continuous** data is quantitative data that needs to be **measured**
 - Continuous data can take **any value** within a range of infinite values
 - For example: the height of a student
- **Age** can be **discrete or continuous** depending on the context or how it is defined
 - If you mean how many years old a person is then this is discrete
 - If you mean how long a person has been alive then this is continuous

What is the difference between a population and a sample?

- The **population** refers to the **whole set** of things which you are interested in
 - For example: if a vet wanted to know how long a typical French bulldog slept for in a day then the population would be all the French bulldogs in the world
- A **sample** refers to a **subset of the population** which is used to collect data from
 - For example: the vet might take a sample of French bulldogs from different cities and record how long they sleep in a day
- A **sampling frame** is a **list** of all members of the **population**
 - For example: a list of employees' names within a company
- Using a **sample instead of a population**:
 - Is quicker and cheaper
 - Leads to less data needing to be analysed
 - Might not fully represent the population
 - Might introduce bias

Sampling Techniques

What is a random sample and a biased sample?

- A **random sample** is where every member of the population has an equal chance of being included in the sample
- A **biased sample** is where the sample is **not random**

What sampling techniques do I need to know?

Simple random sampling

- **Simple random sampling** is where every group of members from the population has an **equal probability** of being selected for the sample
- To carry this out you would...
 - uniquely number every member of a population
 - randomly select n different numbers using a random number generator or a form of lottery (where numbers are selected randomly)
- **Effectiveness:**
 - Useful when you have a small population or want a small sample (such as children in a class)
 - It can be time-consuming if the sample or population is large
 - This can not be used if it is not possible to number or list all the members of the population (such as fish in a lake)

Systematic sampling

- **Systematic sampling** is where a sample is formed by choosing members of a population at regular intervals using a list
- To carry this out you would...
 - calculate the size of the interval $k = \frac{\text{size of population } (N)}{\text{size of sample } (n)}$
 - choose a random starting point between 1 and k
 - select every k th member after the first one
- **Effectiveness:**
 - Useful when there is a natural order (such as a list of names or a conveyor belt of items)
 - Quick and easy to use
 - This can not be used if it is not possible to number or list all the members of the population (such as penguins in Antarctica)

Stratified sampling

- **Stratified sampling** is where the population is divided into disjoint groups (called strata) and then a random sample is taken from each group (stratum)
- The proportion of a stratum that is sampled is equal to the proportion of the population that belong to that stratum
- To carry this out you would...
 - Calculate the number of members sampled from each stratum

YOUR NOTES



- $\frac{\text{size of sample } (n)}{\text{size of population } (N)} \times \text{number of members in the stratum}$
- Take a random sample from each stratum
- **Effectiveness:**
 - Useful when there are very different groups of members within a population
 - The sample will be representative of the population structure
 - The members selected from each stratum are chosen randomly
 - This can not be used if the population can not be split into groups or if the groups overlap

YOUR NOTES



Quota sampling

- **Quota sampling** is where the population is split into groups (like stratified sampling) and members of the population are selected until each quota is filled
- To carry this out you would...
 - Calculate how many people you need from each group
 - Select members from each group until that quota is filled
 - The members do not have to be selected randomly
- **Effectiveness:**
 - Useful when collecting data by asking people who walk past you in a public place or when a sampling frame is not available
 - This can introduce bias as some members of the population might choose not to be included in the sample

Convenience sampling

- **Convenience sampling** is where a sample is formed using available members of the population who fit the criteria
- To carry this out you would...
 - Select members that are easiest to reach
- **Effectiveness:**
 - Useful when a list of the population is not possible
 - This is unlikely to be representative of the population structure
 - This is likely to produce biased results

What are the main criticisms of sampling techniques?

- Most sampling techniques can be improved by taking a larger sample
- Sampling can introduce bias - so you want to minimise the bias within a sample
 - To minimise bias the sample should be as close to random as possible
- A sample only gives information about those members
 - Different samples may lead to different conclusions about the population



Worked Example

Mike is a biologist studying mice in an open enclosure. He has access to approximately 540 field mice and 260 harvest mice. Mike wants to sample 10 mice and he wants the proportions of the two types of mice in his sample to reflect their respective proportions of the population.

a)

Calculate the number of field mice and harvest mice that Mike should include in his sample.

Total number of mice
 $540 + 260 = 800$

Field mice $\frac{540}{800} \times 10 = 6.75$

Harvest mice $\frac{260}{800} \times 10 = 3.25$

Fraction of field mice
 Sample size
 Fraction of harvest mice

Include 7 field mice and 3 harvest mice

b)

Given that Mike does not have a list of all mice in the enclosure, state the name of this sampling method.

No list of population so can not be a random sample

Quota sampling

c)

Suggest one way in which Mike could improve his sampling method.

Mark could improve his sampling method by increasing his sample size

Reliability of Data

How can I decide if data is reliable?

- Data from a sample is reliable if similar results would be obtained from a different sample from the same population
- The sample should be **representative** of the population
- The sample should be **big enough**
 - Sampling a small proportion of a population is unlikely to be reliable

What can cause data to be unreliable?

- If the sample is **biased**
 - It is **not random**
- If **errors** are made when collecting data
 - Numbers could be recorded incorrectly, duplicated or missed out
- If the person collecting the data **favours some members** over others
 - They might seek out members who will lead to a desired outcome
 - They might exclude members if they would cause the sample to oppose the desired outcome
- If a significant proportion of **data is missing**
 - Some data may be unavailable
 - Some members might decide not to be part of the sample
 - This will mean the results are not necessarily representative of the population

YOUR NOTES



4.1.2 Statistical Measures

YOUR NOTES



Mean, Mode, Median

What are the mean, mode and median?

- Mean, median and mode are **measures of central tendency**
 - They describe where the centre of the data is
- They are all types of **averages**
- In statistics it is important to be specific about which average you are referring to
- The **units** for the mean, mode and median are the **same** as the units for the data

How are the mean, mode, and median calculated for ungrouped data?

- The **mode** is the value that occurs **most often** in a data set
 - It is possible for there to be **more than one mode**
 - It is possible for there to be **no mode**
 - In this case **do not** say the mode is zero
- The **median** is the **middle** value when the data is in **order of size**
 - If there are two values in the middle then the median is the **midpoint** of the two values
- The **mean** is the **sum** of all the values **divided by the number of values**

$$\bar{x} = \frac{1}{n} \sum_{i=1}^n x_i$$

- Where $\sum_{i=1}^n x_i = x_1 + x_2 + \dots + x_n$ is the sum of the n pieces of data
- The mean can be represented by the symbol μ
- Your **GDC** can calculate these statistical measures if you input the data using the statistics mode



Worked Example

Find the mode, median and mode for the data set given below.

43 29 70 51 64 43

Mode is the most common

$$\text{Mode} = 43$$

Median is the middle when in order

29 43 43 51 64 70

$$\begin{array}{c} \uparrow \\ \frac{43+51}{2} = 47 \end{array}$$

$$\text{Median} = 47$$

$$\text{Mean} = \frac{\sum x}{n}$$

$$\sum x = 300 \text{ and } n = 6 \quad \frac{300}{6} = 50$$

$$\text{Mean} = 50$$

Quartiles & Range

YOUR NOTES



What are quartiles?

- **Quartiles** are **measures of location**
- Quartiles divide a population or data set into **four equal sections**
 - The **lower quartile, Q_1** splits the lowest 25% from the highest 75%
 - The **median, Q_2** splits the lowest 50% from the highest 50%
 - The **upper quartile, Q_3** splits the lowest 75% from the highest 25%
- There are different methods for finding quartiles
 - Values obtained by hand and using technology may differ
- You will be expected to use your GDC to calculate the quartiles

What are the range and interquartile range?

- The **range** and **interquartile range** are both **measures of dispersion**
 - They describe how spread out the data is
- The **range** is the largest value of the data minus the smallest value of the data
- The **interquartile range** is the range of the central 50% of data
 - It is the upper quartile minus the lower quartile

$$\text{IQR} = Q_3 - Q_1$$

- This is given in the **formula booklet**
- The **units** for the range and interquartile range are the **same** as the units for the data



Worked Example

Find the range and interquartile range for the data set given below.

43 29 70 51 64 43

$$\text{Range} = \text{Maximum} - \text{Minimum}$$

$$70 - 29$$

$$\text{Range} = 41$$

Find upper and lower quartiles using GDC


$$Q_1 = 43 \quad \text{and} \quad Q_3 = 64$$

$$\text{IQR} = Q_3 - Q_1$$

$$64 - 43$$

$$\text{IQR} = 21$$

By hand





Standard Deviation & Variance

What are the standard deviation and variance?

- The **standard deviation** and **variance** are both **measures of dispersion**
 - They describe how spread out the data is in relation to the mean
- The **variance** is the **mean** of the **squares** of the **differences** between **the values and the mean**
 - Variance is denoted σ^2
- The **standard deviation** is the **square-root** of the **variance**
 - Standard deviation is denoted σ
- The **units** for the standard deviation are the **same** as the units for the data
- The **units** for the variance are the **square** of the units for the data

How are the standard deviation and variance calculated for ungrouped data?

- In the exam you will be expected to use the statistics function on your **GDC** to calculate the standard deviation and the variance
- Calculating the standard deviation and the variance by hand may deepen your understanding

- The formula for **variance** is $\sigma^2 = \frac{1}{n} \sum_{i=1}^n (x_i - \bar{x})^2$

- This can be rewritten as

$$\sigma^2 = \frac{1}{n} \sum_{i=1}^n x_i^2 - \bar{x}^2$$

- The formula for **standard deviation** is $\sigma = \sqrt{\frac{1}{n} \sum_{i=1}^n (x_i - \bar{x})^2}$

- This can be rewritten as

$$\sigma = \sqrt{\frac{1}{n} \sum_{i=1}^n x_i^2 - \bar{x}^2}$$

- You **do not** need to learn these formulae as you will use your GDC to calculate these



Worked Example

Find the variance and standard deviation for the data set given below.

43 29 70 51 64 43

Find variance and standard deviation using GDC

$$\sigma_x^2 = 189.333... \quad \text{and} \quad \sigma_x = 13.759...$$

$$\text{Variance} = 189 \text{ (3sf)}$$

$$\text{Standard deviation} = 13.8 \text{ (3sf)}$$

By hand

$$\sigma^2 = \frac{\sum x^2}{n} - \bar{x}^2$$

$$\sum x^2 = 16136 \quad \bar{x} = 50 \quad n = 6$$

$$\sigma^2 = \frac{16136}{6} - 50^2 = 189.333...$$

$$\sigma = \sqrt{189.333...} = 13.759...$$

YOUR NOTES



4.1.3 Frequency Tables

YOUR NOTES



Ungrouped Data

How are frequency tables used for ungrouped data?

- Frequency tables can be used for ungrouped data when you have lots of the same values within a data set
 - They can be used to collect and present data easily
- If the value 4 has a frequency of 3 this means that there are three 4's in the data set

How are measures of central tendency calculated from frequency tables with ungrouped data?

- The **mode** is the value that has the **highest frequency**
- The **median** is the **middle** value
 - Use cumulative frequencies (running totals) to find the median
- The **mean** can be calculated by
 - Multiplying each value x_i by its frequency f_i
 - Summing to get $\sum f_i x_i$
 - Dividing by the total frequency $n = \sum f_i$
 - This is given in the formula booklet

$$\bar{x} = \frac{\sum_{i=1}^k f_i x_i}{n}$$

- Your **GDC** can calculate these statistical measures if you input the values and their frequencies using the statistics mode

How are measures of dispersion calculated from frequency tables with ungrouped data?

- The **range** is the largest value of the data minus the smallest value of the data
- The **interquartile range** is calculated by

$$\text{IQR} = Q_3 - Q_1$$

- The **quartiles** can be found by using your GDC and inputting the values and their frequencies
- The **standard deviation** and **variance** can be calculated by hand using the formulae
 - **Variance**

$$\sigma^2 = \frac{1}{n} \sum_{i=1}^k f_i x_i^2 - \bar{x}^2$$

- **Standard deviation**

$$\sigma = \sqrt{\frac{1}{n} \sum_{i=1}^k f_i X_i^2 - \bar{X}^2}$$

- You **do not need to learn** these formulae as you will be expected to use your GDC to find the standard deviation and variance
 - You may want to see these formulae to deepen your understanding



Exam Tip

- Always check whether your answers make sense when using your GDC
 - The value for a measure of central tendency should be within the range of data

YOUR NOTES





Worked Example

The frequency table below gives information number of pets owned by 30 students in a class.

Number of pets	0	1	2	3
Frequency	11	5	8	6

Find

a)
the mode.

Mode = value with highest frequency

$$\text{Mode} = 0$$

b)
the median.

Median = middle value

$n = 30$ so median is midpoint of 15th and 16th

Number of pets	0	1	2	3
Cumulative frequency	11	16	24	30

$$\text{Median} = 1$$

c)
the mean.

Formula
Booklet

Mean, \bar{x} , of a set of data	$\bar{x} = \frac{\sum_{i=1}^k f_i x_i}{n}$	$n = \sum_{i=1}^k f_i$
------------------------------------	--	------------------------

$$\bar{x} = \frac{\sum f_i x_i}{n} = \frac{11 \times 0 + 5 \times 1 + 8 \times 2 + 6 \times 3}{11 + 5 + 8 + 6} = \frac{39}{30}$$

$$\text{Mean} = 1.3$$

d)
the standard deviation.

Use GDC $\sigma_x = 1.159\dots$

Standard deviation = 1.16 (3sf)

YOUR NOTES



Grouped Data

How are frequency tables used for grouped data?

- Frequency tables can be used for grouped data when you have lots of the same values within the same interval
 - Class intervals will be written using inequalities and without gaps
 - $10 \leq x < 20$ and $20 \leq x < 30$
 - If the class interval $10 \leq x < 20$ has a frequency of 3 this means there are three values in that interval
 - You do not know the **exact data values** when you are given grouped data

How are measures of central tendency calculated from frequency tables with grouped data?

- The **modal class** is the class that has the **highest frequency**
 - This is for equal class intervals only
- The **median** is the **middle** value
 - The exact value can not be calculated but it can be estimated by using a **cumulative frequency graph**
- The **exact mean** can not be calculated as you do not have the raw data
- The **mean** can be **estimated** by
 - Identifying the mid-interval value (midpoint) x_i for each class
 - Multiplying each value by the class frequency f_i
 - Summing to get $\sum f_i x_i$
 - Dividing by the total frequency $n = \sum f_i$
 - This is given in the formula booklet

$$\bar{x} = \frac{\sum_{i=1}^k f_i x_i}{n}$$

- Your **GDC** can estimate the mean if you input the mid-interval values and the class frequencies using the statistics mode

How are measures of dispersion calculated from frequency tables with grouped data?

- The exact **range** can not be calculated as the largest and smallest values are unknown
- The **interquartile range** can be estimated by

$$\text{IQR} = Q_3 - Q_1$$

- **Estimates** of the **quartiles** can be found by using a **cumulative frequency graph**
- The **standard deviation** and **variance** can be estimated using the mid-interval values x_i in the formulae
 - **Variance**

$$\sigma^2 = \frac{1}{n} \sum_{i=1}^k f_i x_i^2 - \bar{x}^2$$

YOUR NOTES



◦ **Standard deviation**

$$\sigma = \sqrt{\frac{1}{n} \sum_{i=1}^k f_i x_i^2 - \bar{x}^2}$$

- You **do not need to learn** these formulae as you will be expected to use your GDC to estimate the standard deviation and variance using the mid-interval values
 - You may want to see these formulae to deepen your understanding

**Exam Tip**

- As you can only estimate statistical measures from a grouped frequency table it is good practice to indicate that the values are not exact
 - You can do this by rounding values rather than leaving as surds and fractions
 - $\bar{x} = 0.333$ (3sf) rather than $\bar{x} = \frac{1}{3}$

YOUR NOTES





Worked Example

The table below shows the heights in cm of a group of 25 students.

Height, h	Frequency
$150 \leq h < 155$	3
$155 \leq h < 160$	5
$160 \leq h < 165$	9
$165 \leq h < 170$	7
$170 \leq h < 175$	1

a)

Write down the modal class.

Modal class = class with highest frequency

Modal class = $160 \leq h < 165$

b)

Write down the mid-interval value of the modal class.

Mid-interval value = $\frac{\text{Upper boundary} + \text{lower boundary}}{2}$

$$\frac{160 + 165}{2}$$

Mid-interval value = 162.5 cm

c)

Calculate an estimate for the mean height.

Use mid-interval values to estimate the mean

Formula
Booklet

Mean, \bar{x} , of a set of data	$\bar{x} = \frac{\sum_{i=1}^n f_i x_i}{n}$	$n = \sum_{i=1}^n f_i$
------------------------------------	--	------------------------

$$\bar{x} = \frac{3 \times 152.5 + 5 \times 157.5 + 9 \times 162.5 + 7 \times 167.5 + 1 \times 172.5}{3 + 5 + 9 + 7 + 1} = \frac{4052.5}{25}$$

Estimated mean = 162.1 cm

4.1.4 Linear Transformations of Data

YOUR NOTES



Linear Transformations of Data

Why are linear transformations of data used?

- Sometimes data might be very large or very small
- You can apply a **linear transformation** to the data to make the values more manageable
 - You may have heard this referred to as:
 - Effects of constant changes
 - Linear coding
- Linear transformations of data can **affect the statistical measures**

How is the mean affected by a linear transformation of data?

- Let \bar{x} be the **mean** of some data
- If you **multiply each value** by a constant **k** then you will need to **multiply the mean by k**
 - Mean is $k\bar{x}$
- If you **add or subtract** a constant **a** from all the **values** then you will need to **add or subtract the constant a to the mean**
 - Mean is $\bar{x} \pm a$

How is the variance and standard deviation affected by a linear transformation of data?

- Let σ^2 be the **variance** of some data
 - σ is the **standard deviation**
- If you **multiply** each value by a constant **k** then you will need to **multiply the variance by k^2**
 - Variance is $k^2\sigma^2$
 - You will need to **multiply the standard deviation** by the **absolute value** of **k**
 - Standard deviation is $|k|\sigma$
 - If you **add or subtract** a constant **a** from all the **values** then the **variance** and the **standard deviation stay the same**
 - Variance is σ^2
 - Standard deviation is σ



Exam Tip

- If you forget these results in an exam then you can look in the HL section of the formula booklet to see them written in a more algebraic way
 - Linear transformation of a single variable

$$E(aX + b) = aE(X) + b$$

$$\text{Var}(aX + b) = a^2\text{Var}(X)$$

- where $E(\dots)$ means the mean and $\text{Var}(\dots)$ means the variance



Worked Example

A teacher marks his students' tests. The raw mean score is 31 marks and the standard deviation is 5 marks. The teacher standardises the score by doubling the raw score and then adding 10.

a)

Calculate the mean standardised score.

If data is multiplied by k then mean is multiplied by k

If k is added to data then k is added to the mean

$$31 \times 2 + 10$$

$$\text{Mean of standardised scores} = 72$$

b)

Calculate the standard deviation of the standardised scores.

If data is multiplied by k then standard deviation is multiplied by $|k|$

If k is added to data then standard deviation is unchanged

$$5 \times 2$$

$$\text{Standard deviation of standardised scores} = 10$$



Outliers

What are outliers?

- Outliers are extreme data values that do not fit with the rest of the data
 - They are either a lot bigger or a lot smaller than the rest of the data
- Outliers are defined as values that are **more than $1.5 \times \text{IQR}$ from the nearest quartile**
 - x is an outlier if $x < Q_1 - 1.5 \times \text{IQR}$ or $x > Q_3 + 1.5 \times \text{IQR}$
- Outliers can have a big effect on some statistical measures

Should I remove outliers?

- The decision to remove outliers will **depend on the context**
- Outliers **should be removed** if they are found to be **errors**
 - The data may have been recorded incorrectly
 - For example: The number 17 may have been recorded as 71 by mistake
- Outliers **should not be removed** if they are a **valid part of the sample**
 - The data may need to be checked to verify that it is not an error
 - For example: The annual salaries of employees of a business might appear to have an outlier but this could be the director's salary



Worked Example

The ages, in years, of a number of children attending a birthday party are given below.

2, 7, 5, 4, 8, 4, 6, 5, 5, 29, 2, 5, 13

a)

Identify any outliers within the data set.

x is an outlier if $x < Q_1 - 1.5 \times IQR$ or $x > Q_3 + 1.5 \times IQR$

Using GDC

$$Q_1 = 4 \quad \text{and} \quad Q_3 = 7.5 \quad \therefore IQR = 3.5$$

$$Q_1 - 1.5 \times IQR = 4 - 1.5 \times 3.5 = -1.25$$

$$Q_3 + 1.5 \times IQR = 7.5 + 1.5 \times 3.5 = 12.75$$

Outliers are 13 and 29

b)

Suggest which value(s) should be removed. Justify your answer.

13 should not be removed as it is a valid age of a child.

29 should be removed as this is an age of an adult.

4.1.6 Univariate Data

YOUR NOTES




Box Plots

Univariate data is data that is in **one variable**.

What is a box plot (box and whisker diagram)?

- A box plot is a graph that clearly shows key statistics from a data set
 - It shows the **median, quartiles, minimum** and **maximum values** and **outliers**
 - It does not show any other individual data items
- The middle 50% of the data will be represented by the box section of the graph and the lower and upper 25% of the data will be represented by each of the whiskers
- Any **outliers** are represented with a **cross** on the **outside of the whiskers**
 - If there is an outlier then the whisker will end at the value before the outlier
- Only one axis is used when graphing a box plot
- It is still important to make sure the axis has a clear, even scale and is labelled with units



What are box plots useful for?

- Box plots can clearly show the shape of the distribution
 - If a box plot is symmetrical about the median then the data could be **normally distributed**
- Box plots are often used for **comparing two sets of data**
 - Two box plots will be drawn next to each other using the same axis
 - They are useful for **comparing data** because it is easy to see the main shape of the distribution of the data from a box plot
 - You can easily compare the medians and interquartile ranges



Exam Tip

- In an exam you can use your GDC to draw a box plot if you have the raw data
 - Your calculator's box plot can also include outliers so this is a good way to check



Worked Example

The distances, in metres, travelled by 15 snails in a one-minute period are recorded and shown below:

0.5, 0.7, 1.0, 1.1, 1.2, 1.2, 1.2, 1.3, 1.4, 1.4, 1.4, 1.4, 1.5, 1.5, 1.5

a)

i)

Find the values of Q_1 , Q_2 and Q_3 .

ii)

Find the interquartile range.

iii)

Identify any outliers.

Using GDC

$$Q_1 = 1.1 \text{ m} \quad Q_2 = 1.3 \text{ m} \quad Q_3 = 1.4 \text{ m}$$

$$IQR = Q_3 - Q_1 = 1.4 - 1.1$$

$$IQR = 0.3 \text{ m}$$


$$Q_1 - 1.5 \times IQR = 1.1 - 1.5 \times 0.3 = 0.65$$

$$Q_3 + 1.5 \times IQR = 1.4 + 1.5 \times 0.3 = 1.85$$

$$0.5 \text{ m is an outlier}$$

b)

Draw a box plot for the data.



Label outlier with a cross

Use next smallest after outlier

Cumulative Frequency Graphs

YOUR NOTES



What is cumulative frequency?

- The cumulative frequency of x is the running total of the frequencies for the values that are less than or equal to x
- For grouped data you use the upper boundary of a class interval to find the cumulative frequency of that class

What is a cumulative frequency graph?

- A cumulative frequency graph is used with data that has been organised into a **grouped frequency** table
- Some coordinates are plotted
 - The x -coordinates are the **upper boundaries** of the class intervals
 - The y -coordinates are the **cumulative frequencies** of that class interval
- The coordinates are then joined together by hand using a **smooth increasing curve**


What are cumulative frequency graphs useful for?

- They can be used to **estimate** statistical measures
 - Draw a **horizontal line** from the y -axis to the curve
 - For the median: draw the line at 50% of the total frequency
 - For the lower quartile: draw the line at 25% of the total frequency
 - For the upper quartile: draw the line at 75% of the total frequency
 - For the p^{th} percentile: draw the line at $p\%$ of the total frequency
 - Draw a **vertical line** down from the curve to the x -axis
 - This **x -value** is the relevant statistical measure
- They can be used to estimate the number of values that are bigger/small than a given value
 - Draw a **vertical line** from the given value on the x -axis to the curve
 - Draw a **horizontal line** from the curve to the y -axis
 - This value is an estimate for how many values are less than or equal to the given value
 - To estimate the number that is greater than the value subtract this number from the total frequency
 - They can be used to **estimate** the **interquartile range** $IQR = Q_3 - Q_1$
 - They can be used to construct a **box plot** for grouped data




Worked Example

The cumulative frequency graph below shows the lengths in cm, I , of 30 puppies in a training group.



a)

Given that the interval $40 \leq I < 45$ was used when collecting data, find the frequency of this class.




$$16 - 8$$

$$\text{Frequency} = 8$$

b)

Use the graph to find an estimate for the interquartile range of the lengths.



$$\frac{1}{4} \times 30 = 7.5 \quad Q_1 = 39.5$$


$$\frac{3}{4} \times 30 = 22.5 \quad Q_3 = 51.4$$

$$IQR = Q_3 - Q_1 = 51.4 - 39.5$$

$$IQR = 11.9 \text{ cm}$$

c)

Estimate the percentage of puppies with length more than 51 cm.



$$30 - 22 = 8 \text{ puppies}$$

longer than 51 cm

$$\frac{8}{30} \times 100\% = 26.666\ldots\%$$

$$26.7\% \text{ (3sf)}$$

YOUR NOTES



Histograms

What is a (frequency) histogram?

- A frequency histogram clearly shows the frequency of class intervals
 - The classes will have **equal class intervals**
 - The **frequency** will be on the y-axis
 - The bar for a class interval will begin at the lower boundary and end at the upper boundary
- A frequency histogram is **similar to a bar chart**
 - A **bar chart** is used for **qualitative or discrete data** and **has gaps** between the bars
 - A **frequency histogram** is used for **continuous data** and **has no gaps** between bars

What are (frequency) histograms useful for?

- They show the **modal class** clearly
- They show the shape of the distribution
 - It is important the class intervals are of equal width
- They can show whether the variable can be modelled by a **normal distribution**
 - If the shape is symmetrical and bell-shaped

YOUR NOTES






Worked Example

The table below and its corresponding histogram show the mass, in kg, of some new born bottlenose dolphins.

Mass, m kg	Frequency
$4 \leq m < 8$	4
$8 \leq m < 12$	15
$12 \leq m < 16$	19
$16 \leq m < 20$	10
$20 \leq m < 24$	6

a)

Draw a frequency histogram to represent the data.



No gaps

b)

Write down the modal class.

Modal class = class with highest frequency

Modal class = $12 \leq m < 16$

4.1.7 Interpreting Data

YOUR NOTES



Interpreting Data

How do I interpret statistical measures?

- The **mode** is useful for **qualitative data**
 - It is not as useful for quantitative data as there is not always a unique mode
- The **mean includes all values**
 - It is affected by outliers
 - A smaller/larger mean is preferable depending on the scenario
 - A smaller mean time for completing a puzzle is better
 - A bigger mean score on a test is better
- The **median is not affected by outliers**
 - It does not use all the values
- The **range gives the full spread** of the all of the data
 - It is affected by outliers
- The **interquartile range gives the spread of the middle 50%** about the median and is not affected by outliers
 - It does not use all the values
 - A bigger IQR means the data is more spread out about the median
 - A smaller IQR means the data is more centred about the median
- The **standard deviation and variance** use all the values to give a measure of the **average spread** of the data about the mean
 - They are affected by outliers
 - A bigger standard deviation means the data is more spread out about the mean
 - A smaller standard deviation means the data is more centred about the mean

How do I choose which diagram to use to represent data?

- **Box plots**
 - Can be used with ungrouped **univariate** data
 - Shows the range, interquartile range and quartiles clearly
 - Very useful for comparing data patterns quickly
- **Cumulative frequency graphs**
 - Can be used with continuous grouped univariate data
 - Shows the running total of the frequencies that fall below the upper bound of each class
- **Histograms**
 - Can be used with continuous grouped univariate data
 - Used with equal class intervals
 - Shows the frequencies of the group
- **Scatter diagrams**
 - Can be used with ungrouped **bivariate** data
 - Shows the graphical relationship between the variables

How do I compare two or more data sets?


- Compare a **measure of central tendency**
 - If the data **contains outliers** - use the **median**



- If the data is **roughly symmetrical** – use the mean
- Compare a **measure of dispersion**
 - If the data **contains outliers** – use the **interquartile range**
 - If the data is **roughly symmetrical** – use the **standard deviation**
- Consider whether it is better to have a smaller or bigger average
 - This will depend on the context
 - A smaller average time for completing a puzzle is better
 - A bigger average score on a test is better
- Consider whether it is better to have a smaller or bigger spread
 - Usually a smaller spread means it is more consistent
- Always relate the **comparisons to the context** and consider reasons
 - Consider the **sampling technique** and the **data collection** method

? Worked Example

The box plots below show the waiting times for the two doctor surgeries, HealthHut and FitFirst.



Compare the two distributions of waiting times in context.

Compare :

- a measure of central tendency
- a measure of dispersion

HealthHut's median waiting time is smaller than FitFirst's ($20 < 24$). On average patients get seen quicker at HealthHut.

FitFirst's interquartile range is smaller than HealthHut's ($13 < 19$). There is less variability of waiting times at FitFirst.

4.2 Correlation & Regression

4.2.1 Bivariate data

Scatter Diagrams

What does bivariate data mean?

- **Bivariate data** is data which is collected on **two variables** and looks at how one of the factors affects the other
 - Each data value from one variable will be **paired** with a data value from the other variable
 - The two variables are often related, but do not have to be

What is a scatter diagram?

- A **scatter diagram** is a way of graphing bivariate data
 - One variable will be on the **x-axis** and the other will be on the **y-axis**
 - The variable that can be **controlled** in the data collection is known as the **independent** or **explanatory variable** and is plotted on the **x-axis**
 - The variable that is **measured** or discovered in the data collection is known as the **dependent** or **response variable** and is plotted on the **y-axis**
- Scatter diagrams can contain **outliers** that do not follow the trend of the data



Exam Tip

- If you use scatter diagrams in your Internal Assessment then be aware that finding outliers for bivariate data is different to finding outliers for univariate data
 - (x, y) could be an outlier for the bivariate data even if x and y are not outliers for their separate univariate data

YOUR NOTES




Correlation

YOUR NOTES



What is correlation?

- **Correlation** is how the **two variables change in relation to each other**
 - Correlation could be the result of a **causal relationship** but this is not always the case
- **Linear correlation** is when the changes are proportional to each other
- **Perfect linear correlation** means that the bivariate data will all lie on a straight line on a scatter diagram
- When describing correlation mention
 - The type of the correlation
 - **Positive correlation** is when an **increase** in one variable results in the other variable **increasing**
 - **Negative correlation** is when an **increase** in one variable results in the other variable **decreasing**
 - **No linear correlation** is when the data points don't appear to follow a trend
 - The strength of the correlation
 - **Strong linear correlation** is when the data points lie **close** to a **straight line**
 - **Weak linear correlation** is when the data points are **not close** to a **straight line**
- If there is **strong linear correlation** you can draw a **line of best fit** (by eye)
 - The line of best fit will pass through the mean point (\bar{x}, \bar{y})
 - If you are asked to draw a line of best fit
 - Plot the mean point
 - Draw a line going through it that follows the trend of the data



Copyright © Save My Exams. All Rights Reserved

What is the difference between correlation and causation?

- It is important to be aware that just because correlation exists, it does not mean that the change in one of the variables is **causing** the change in the other variable

- **Correlation does not imply causation!**
- If a change in one variable **causes** a change in the other then the two variables are said to have a **causal relationship**
 - Observing correlation between two variables does **not always** mean that there is a causal relationship
 - There could be **underlying factors** which is causing the correlation
 - Look at the two variables in question and consider the context of the question to decide if there could be a causal relationship
 - If the two variables are temperature and number of ice creams sold at a park then it is likely to be a causal relationship
 - Correlation may exist between global temperatures and the number of monkeys kept as pets in the UK but they are unlikely to have a causal relationship

YOUR NOTES






Worked Example

A teacher is interested in the relationship between the number of hours her students spend on a phone per day and the number of hours they spend on a computer. She takes a sample of nine students and records the results in the table below.

Hours spent on a phone per day	7.6	7.0	8.9	3.0	3.0	7.5	2.1	1.3	5.8
Hours spent on a computer per day	1.7	1.1	0.7	5.8	5.2	1.7	6.9	7.1	3.3

- a)
Draw a scatter diagram for the data.




- b)
Describe the correlation.

Strong negative linear correlation

- c)
Draw a line of best fit.

Mean point $(\bar{x}, \bar{y}) = (5.133..., 3.722...)$



Plot the mean point

Draw it by eye

YOUR NOTES











4.2.2 Correlation Coefficients

YOUR NOTES


PMCC
What is Pearson's product-moment correlation coefficient?

- Pearson's product-moment correlation coefficient (PMCC) is a way of giving a numerical value to a **linear relationship** of bivariate data
- The PMCC of a sample is denoted by the letter r
 - r can take any value such that $-1 \leq r \leq 1$
 - A **positive value** of r describes **positive correlation**
 - A **negative value** of r describes **negative correlation**
 - $r = 0$ means there is **no linear correlation**
 - $r = 1$ means **perfect positive linear** correlation
 - $r = -1$ means **perfect negative linear** correlation
 - The closer to 1 or -1 the stronger the correlation



Copyright © Save My Exams. All Rights Reserved

How do I calculate Pearson's product-moment correlation coefficient (PMCC)?

- You will be expected to use the statistics mode on your GDC to calculate the PMCC
- The formula can be useful to deepen your understanding

$$r = \frac{S_{xy}}{S_x S_y}$$



- $S_{xy} = \sum_{i=1}^n x_i y_i - \frac{1}{n} \left(\sum_{i=1}^n x_i \right) \left(\sum_{i=1}^n y_i \right)$ is linked to the **covariance**
- $S_x = \sqrt{\sum_{i=1}^n x_i^2 - \frac{1}{n} \left(\sum_{i=1}^n x_i \right)^2}$ and $S_y = \sqrt{\sum_{i=1}^n y_i^2 - \frac{1}{n} \left(\sum_{i=1}^n y_i \right)^2}$ are linked to the **variances**
 - You **do not need to learn this** as using your GDC will be expected

When does the PMCC suggest there is a linear relationship?

- **Critical values** of r indicate when the PMCC would suggest there is a linear relationship
 - In your exam you will be given critical values where appropriate
 - Critical values will depend on the size of the sample
- If the **absolute value** of the **PMCC** is **bigger** than the **critical value** then this suggests a linear model is appropriate

Spearman's Rank

What is Spearman's rank correlation coefficient?

- Each data is ranked from biggest to smallest
 - For n data values they are ranked from 1 to n
 - 1 for the biggest x -value and 1 for the biggest y -value
- Spearman's rank correlation coefficient is a way to measure linear correlation between the rankings of the data
- Spearman's rank of a sample is denoted by r_s
 - r_s can take any value such that $-1 \leq r_s \leq 1$
 - A **positive value** of r_s describes a **degree of agreement** between the rankings
 - A **negative value** of r_s describes a **degree of disagreement** between the rankings
 - $r_s = 0$ means there is **no correlation** between the rankings
 - $r_s = 1$ means the rankings are in **complete agreement**
 - An increase in one variable means an increase in the other
 - $r_s = -1$ means the rankings are in **complete disagreement**
 - An increase in one variable means a decrease in the other
 - If $r_s = 1$ or $r_s = -1$ then the data is said to have a **monotonic** relationship
 - This means either the points are always increasing or always decreasing
 - The closer to 1 or -1 the stronger the correlation of the rankings

How do I calculate Spearman's rank correlation coefficient (PMCC)?

- Rank each set of data independently
 - 1 to n for the x -values
 - 1 to n for the y -values
- If some values are equal then give each the average of the ranks they would occupy
 - For example: if the 3rd, 4th and 5th highest values are equal then give each the ranking of 4
 - $\frac{3 + 4 + 5}{3} = 4$
- Calculate the PMCC of the **rankings** using your GDC
 - This value is **Spearman's rank correlation coefficient**

Appropriateness & Limitations

Which correlation coefficient should I use?

- **Pearson's PMCC** tests for a **linear relationship** between two variables
 - It will not tell you if the variables have a non-linear relationship
 - Such as exponential growth
 - Use this if you are interested in a linear relationship
- **Spearman's rank** tests for a **monotonic relationship** (always increasing or always decreasing) between two variables
 - It will not tell you what function can be used to model the relationship
 - Both linear relationships and exponential relationships can be monotonic
 - Use this if you think there is a non-linear monotonic relationship

How are Pearson's and Spearman's correlation coefficients connected?

- If there is **linear correlation** then the relationship is also **monotonic**
 - $r = 1 \Rightarrow r_s = 1$
 - $r = -1 \Rightarrow r_s = -1$
 - However the **converse is not true**
- It is possible for Spearman's rank to be 1 (or -1) but for the PMCC to be different
 - For example: data that follows an **exponential growth model**
 - $r_s = 1$ as the points are always increasing
 - $r < 1$ as the points do not lie on a straight line

Are Pearson's and Spearman's correlation coefficients affected by outliers?

- Pearson's PMCC **is** affected by outliers
 - as it uses the numerical value of each data point
- Spearman's rank is **not usually** affected by outliers
 - as it only uses the ranks of each data point



Exam Tip

- You can use your GDC to plot the scatter diagram to help you visualise the data

YOUR NOTES





Worked Example

The table below shows the scores of eight students for a maths test and an English test.

Maths (x)	7	18	37	52	61	68	75	82
English (y)	5	3	9	12	17	41	49	97

a)

Write down the value of Pearson's product-moment correlation coefficient, r .

Enter data into GDC.

$$r = 0.79433\dots$$

$$r = 0.794 \text{ (3sf)}$$

b)

Find the value of Spearman's rank correlation coefficient, r_s .

Rank the data

x rank	8	7	6	5	4	3	2	1
y rank	7	8	6	5	4	3	2	1

Find PMCC of ranks

$$r_s = 0.97619\dots$$

$$r_s = 0.976 \text{ (3sf)}$$

c)

Comment on the values of the two correlation coefficients.

The value of r suggests there is strong positive linear correlation. The value of r_s suggests that the relationship is almost monotonic.

4.2.3 Linear Regression

YOUR NOTES



Linear Regression

What is linear regression?

- If **strong linear correlation** exists on a scatter diagram then the data can be modelled by a **linear model**
 - Drawing lines of best fit by eye is not the best method as it can be difficult to judge the best position for the line
- The **least squares regression line** is the line of best fit that minimises the **sum of the squares** of the gap between the line and each data value
 - This is usually called the **regression line of y on x**
 - It can be calculated by looking at the vertical distances between the line and the data values
- The **regression line of y on x** is written in the form $y = ax + b$
- a is the **gradient** of the line
 - It represents the change in y for each individual unit change in x
 - If a is **positive** this means y **increases** by a for a unit increase in x
 - If a is **negative** this means y **decreases** by $|a|$ for a unit increase in x
- b is the **y – intercept**
 - It shows the value of y when x is zero
- You are expected to use your **GDC** to find the equation of the regression line
 - Enter the bivariate data and choose the **model “ $ax + b$ ”**
 - Remember the **mean point** (\bar{x}, \bar{y}) will lie on the regression line

How do I use a regression line?

- The equation of the regression line can be used to decide what type of correlation there is if there is no scatter diagram
 - If a is **positive** then the data set has **positive correlation**
 - If a is **negative** then the data set has **negative correlation**
- The equation of the regression line can also be used to **predict** the value of a **dependent variable (y)** from an **independent variable (x)**
 - The equation should **only be used** to make **predictions for y**
 - Using a y on x line to **predict x is not always reliable**
 - Making a prediction **within the range** of the given data is called **interpolation**
 - This is usually reliable
 - The stronger the correlation the more reliable the prediction
 - Making a prediction **outside of the range** of the given data is called **extrapolation**
 - This is much less reliable
 - The prediction will be more reliable if the number of data values in the original sample set is bigger



Exam Tip

- Once you calculate the values of a and b store them in your GDC
 - This means you can use the full display values rather than the rounded values when using the linear regression equation to predict values
 - This avoids rounding errors

YOUR NOTES





Worked Example

Barry is a music teacher. For 7 students, he records the time they spend practising per week (x hours) and their score in a test (y %).

Time (x)	2	5	6	7	10	11	12
Score (y)	11	49	55	75	63	68	82

a)

Write down the equation of the regression line of y on x , giving your answer in the form $y = ax + b$ where a and b are constants to be found.

Enter data into GDC

a is the coefficient of x $a = 5.5680\dots$

b is the constant term $b = 15.4136\dots$

$$y = 5.57x + 15.4$$

b)

Give an interpretation of the value of a .

$a = 5.57$ means that the model suggests that the score increases by 5.57% for every extra hour of practice.

c)

Another of Barry's students practises for 15 hours a week, estimate their score. Comment on the validity of this prediction.

Substitute $x = 15$

$$y = (5.5680\dots) \times 15 + (15.4136\dots) = 98.93\dots$$

The model predicts a score of 98.9% but this is unreliable as $x = 15$ is outside the range of data. Therefore extrapolation is being used.

4.3 Probability

4.3.1 Probability & Types of Events

YOUR NOTES



Probability Basics

What key words and terminology are used with probability?

- An **experiment** is a repeatable activity that has a result that can be observed or recorded
 - **Trials** are what we call the repeats of the experiment
- An **outcome** is a possible result of a trial
- An **event** is an outcome or a collection of outcomes
 - Events are usually denoted with capital letters: A , B , etc
 - $n(A)$ is the number of outcomes that are included in event A
 - An event can have one or more than one outcome
- A **sample space** is the set of all possible outcomes of an experiment
 - This is denoted by U
 - $n(U)$ is the total number of outcomes
 - It can be represented as a **list** or a **table**

How do I calculate basic probabilities?

- If all outcomes are **equally likely** then probability for each outcome is the same
 - Probability for each outcome is $\frac{1}{n(U)}$
- **Theoretical probability** of an event can be calculated without using an experiment by dividing the number of outcomes of that event by the total number of outcomes

$$P(A) = \frac{n(A)}{n(U)}$$

- This is given in the **formula booklet**
- Identifying all possible outcomes either as a list or a table can help
- **Experimental probability** (also known as **relative frequency**) of an outcome can be calculated using results from an experiment by dividing its frequency by the number of trials
 - **Relative frequency** of an outcome is $\frac{\text{Frequency of that outcome from the trials}}{\text{Total number of trials } (n)}$

How do I calculate the expected number of occurrences of an outcome?

- **Theoretical probability** can be used to calculate the **expected number of occurrences** of an outcome from n trials
- If the probability of an outcome is p and there are n trials then:
 - The expected number of occurrences is **np**
 - This **does not mean** that there will **exactly np occurrences**
 - If the experiment is repeated multiple times then we expect the number of occurrences to average out to be np

What is the complement of an event?

- The probabilities of all the outcomes **add up to 1**

- Complementary events are when there are **two events** and **exactly one** of them will occur
 - One event has to occur but both events can not occur at the same time
- The **complement of event A** is the event where event **A does not happen**
 - This can be thought of as **not A**
 - This is denoted A'

$$P(A) + P(A') = 1$$

- This is in the **formula booklet**
- It is commonly written as $P(A') = 1 - P(A)$

What are different types of combined events?

- The **intersection** of two events (A and B) is the event where **both A and B** occur
 - This can be thought of as **A and B**
 - This is denoted as $A \cap B$
- The **union** of two events (A and B) is the event where **A or B or both occur**
 - This can be thought of as **A or B**
 - This is denoted $A \cup B$
- The event where A occurs given that event B has occurred is called **conditional probability**
 - This can be thought as **A given B**
 - This is denoted $A|B$

How do I find the probability of combined events?

- The probability of A or B (or both) occurring can be found using the formula

$$P(A \cup B) = P(A) + P(B) - P(A \cap B)$$

- This is given in the **formula booklet**
 - You subtract the probability of A and B both occurring because it has been included twice (once in $P(A)$ and once in $P(B)$)
- The probability of A and B occurring can be found using the formula

$$P(A \cap B) = P(A)P(B|A)$$

- A rearranged version is given in the **formula booklet**
- Basically you multiply the probability of A by the probability of B then happening



Exam Tip

- In an exam drawing a Venn diagram or tree diagram can help even if the question does not ask you to

YOUR NOTES





Worked Example

Dave has two fair spinners, A and B . Spinner A has three sides numbered 1, 4, 9 and spinner B has four sides numbered 2, 3, 5, 7. Dave spins both spinners and forms a two-digit number by using the spinner A for the first digit and spinner B for the second digit.

T is the event that the two-digit number is a multiple of 3.

a)

List all the possible two-digit numbers.

A two-way table would be a systematic way to list all the outcomes

	2	3	5	7
1	12	13	15	17
4	42	43	45	47
9	92	93	95	97

b)

Find $P(T)$.

$$P(T) = \frac{n(T)}{n(U)} \leftarrow \begin{array}{l} \text{Number of multiples of 3} \\ \text{Total number of outcomes} \end{array}$$

{12, 15, 42, 45, 93} are the multiples of 3

$$P(T) = \frac{5}{12}$$

c)

Find $P(T')$.

$$P(T) + P(T') = 1 \Rightarrow P(T') = 1 - P(T)$$

$$P(T') = 1 - \frac{5}{12}$$

$$P(T') = \frac{7}{12}$$



Independent & Mutually Exclusive Events

What are mutually exclusive events?

- Two events are **mutually exclusive** if they **can not both happen at once**
 - For example: when rolling a dice the events “getting a prime number” and “getting a 6” are mutually exclusive
- If A and B are mutually exclusive events then:
 - $P(A \cap B) = 0$
- **Complementary events** are mutually exclusive

What are independent events?

- Two events are **independent** if **one occurring does not affect the probability of the other occurring**
 - For example: when flipping a coin twice the events “getting a tails on the first flip” and “getting a tails on the second flip” are independent
- If A and B are independent events then:
 - $P(A|B) = P(A)$ and $P(B|A) = P(B)$
- If A and B are independent events then:
 - $P(A \cap B) = P(A)P(B)$
 - This is given in the **formula booklet**
 - This is a useful formula to test whether two events are statistically independent

How do I find the probability of combined mutually exclusive events?

- If A and B are **mutually exclusive** events then

$$P(A \cup B) = P(A) + P(B)$$

- This is given in the **formula booklet**
 - This occurs because $P(A \cap B) = 0$
- For any two events A and B the events $A \cap B$ and $A \cap B'$ are **mutually exclusive** and A is the **union** of these two events
 - $P(A) = P(A \cap B) + P(A \cap B')$
 - This works for any two events A and B



? Worked Example

a)

A student is chosen at random from a class. The probability that they have a dog is 0.8, the probability they have a cat is 0.6 and the probability that they have a cat or a dog is 0.9.

Find the probability that the student has both a dog and a cat.

Let D be event "has a dog" and C be "has a cat"

$$P(D \cup C) = P(D) + P(C) - P(D \cap C)$$

$$0.9 = 0.8 + 0.6 - P(D \cap C)$$

$$P(D \cap C) = 0.5$$

b)

Two events, Q and R , are such that $P(Q) = 0.8$ and $P(Q \cap R) = 0.1$.

Given that Q and R are independent, find $P(R)$.

$$Q \text{ and } R \text{ independent} \Rightarrow P(Q \cap R) = P(Q)P(R)$$

$$0.1 = 0.8 \times P(R) \quad \therefore P(R) = \frac{0.1}{0.8}$$

$$P(R) = 0.125 \text{ or } \frac{1}{8}$$

c)

Two events, S and T , are such that $P(S) = 2P(T)$.

Given that S and T are mutually exclusive and that $P(S \cup T) = 0.6$ find $P(S)$ and $P(T)$.

$$S \text{ and } T \text{ mutually exclusive} \Rightarrow P(S \cup T) = P(S) + P(T)$$

$$0.6 = P(S) + P(T)$$

$$0.6 = 2P(T) + P(T) \quad P(S) = 2P(T)$$

$$0.6 = 3P(T)$$

$$P(T) = 0.2 \text{ and } P(S) = 0.4$$

4.3.2 Conditional Probability

YOUR NOTES



Conditional Probability

What is conditional probability?

- **Conditional probability** is where the probability of an **event** happening can vary depending on the outcome of a prior event
- The event A happening **given that** event B has happened is denoted $A|B$
- A common example of conditional probability involves selecting multiple objects from a bag **without replacement**
 - The probability of selecting a certain item changes depending on what was selected before
 - This is because the total number of items will change as they are not replaced once they have been selected

How do I calculate conditional probabilities?

- Some conditional probabilities can be calculated by using counting outcomes
 - Probabilities without replacement can be calculated like this
 - For example: There are 10 balls in a bag, 6 of them are red, two of them are selected without replacement
 - To find the probability that the second ball selected is red given that the first one is red count how many balls are left:
 - A red one has already been selected so there are 9 balls left and 5 are red so the probability is $\frac{5}{9}$
- You can use sample space diagrams to find the probability of A given B :
 - reduce your sample space to just include outcomes for event B
 - find the proportion that also contains outcomes for event A
- There is a formula for conditional probability that you can use
 - $$P(A|B) = \frac{P(A \cap B)}{P(B)}$$
 - This is given in the **formula booklet**
 - This can be rearranged to give $P(A \cap B) = P(B)P(A|B)$



Worked Example

In a class of 30 students: 19 students have a dog, 17 students have a cat and 11 have both a dog and a cat. One student is selected at random.

a)

Find the probability that the student has a dog.

Let D be event "has a dog" and C be "has a cat"

$$P(D) = \frac{n(D)}{n(U)} \leftarrow \begin{array}{l} \text{Number who have dogs} \\ \text{Total number of students} \end{array}$$

$$P(D) = \frac{19}{30}$$

b)

Find the probability that the student has a dog given that they have a cat.

17 have a cat of which 11 also have a dog

$$P(D|C) = \frac{11}{17} \quad \text{Could also use } P(D|C) = \frac{P(D \cap C)}{P(C)}$$

c)

Find the probability that the student has a cat given that they have a dog.

19 have a dog of which 11 also have a cat

$$P(C|D) = \frac{11}{19} \quad \text{Could also use } P(C|D) = \frac{P(C \cap D)}{P(D)}$$

4.3.3 Sample Space Diagrams

YOUR NOTES




Venn Diagrams



What is a Venn diagram?

- A Venn diagram is a way to illustrate **events** from an **experiment** and are particularly useful when there is an overlap between possible **outcomes**
- A Venn diagram consists of
 - a **rectangle** representing the **sample space (U)**
 - The rectangle is labelled U
 - Some mathematicians instead use S or ξ
 - a **circle** for each **event**
 - Circles may or may not overlap depending on which **outcomes** are shared between **events**
- The numbers in the circles represent either the **frequency** of that event or the **probability** of that event
 - If the **frequencies** are used then they should **add up to the total frequency**
 - If the **probabilities** are used then they should **add up to 1**

What do the different regions mean on a Venn diagram?

- A' is represented by the regions that are **not in** the A circle
- $A \cap B$ is represented by the region where the A and B circles **overlap**
- $A \cup B$ is represented by the regions that **are in** A or B or both
- Venn diagrams show '**AND**' and '**OR**' statements easily
- Venn diagrams also instantly show **mutually exclusive** events as these circles will **not overlap**
- **Independent** events can not be instantly seen
 - You need to use probabilities to deduce if two events are independent






Copyright © Save My Exams. All Rights Reserved

How do I solve probability problems involving Venn diagrams?

- Draw, or add to a given Venn diagram, filling in as many values as possible from the information provided in the question
- It is usually helpful to work from the centre outwards
 - Fill in **intersections** (overlaps) first
- If two events are independent you can use the formula
 - $P(A \cap B) = P(A)P(B)$
- To find the conditional probability $P(A|B)$
 - Add together the frequencies/probabilities in the B circle
 - This is your denominator
 - Out of those frequencies/probabilities add together the ones that are also in the A circle
 - This is your numerator
 - Evaluate the fraction



Event $A|B$
"A given B"

$$P(A|B) = \frac{P(A \cap B)}{P(B)}$$

← Shade second

← Shade first

$P(A|B) = \frac{\text{"double shading"}}{\text{"single shading"}}$

Copyright © Save My Exams. All Rights Reserved



Exam Tip

- If you struggle to fill in a Venn diagram in an exam:
 - Label the missing parts using algebra
 - Form equations using known facts such as:
 - the sum of the probabilities should be 1
 - $P(A \cap B) = P(A)P(B)$ if A and B are independent events

YOUR NOTES






? Worked Example

40 people are asked if they have sugar and/or milk in their coffee. 21 people have sugar, 25 people have milk and 7 people have neither.

a)

Draw a Venn diagram to represent the information.


Find the centre first



Total should be 40

$$(21-x) + x + (25-x) + 7 = 40$$

$$53 - x = 40 \quad \therefore x = 13$$



b)

One of the 40 people are randomly selected, find the probability that they have sugar but not milk with their coffee.

S and not M is the part of S circle that does not include M

$$P(S \cap M') = \frac{8}{40}$$

Remember to write as a fraction of the total

$$P(S \cap M') = \frac{1}{5}$$

c)

Given that a person who has sugar is selected at random, find the probability that they have milk with their coffee.

Given that sugar has been selected we only want the S circle as our total.

Out of the S circle 13 also have milk

$$P(M|S) = \frac{13}{21}$$

Tree Diagrams

YOUR NOTES




What is a tree diagram?

- A **tree diagram** is another way to show the outcomes of combined events
 - They are very useful for intersections of events
- The events on the branches must be **mutually exclusive**
 - Usually they are an event and its complement
- The probabilities on the second sets of branches **can depend** on the outcome of the first event
 - These are **conditional probabilities**
- When selecting the items from a bag:
 - The second set of branches will be the **same** as the first if the items **are replaced**
 - The second set of branches will be the **different** to the first if the items **are not replaced**

How are probabilities calculated using a tree diagram?

- To find the probability that two events happen together you **multiply** the corresponding probabilities on their branches
 - It is helpful to find the probability of all combined outcomes once you have drawn the tree
- To find the probability of an event you can:
 - **add together** the probabilities of the **combined outcomes** that are part of that event
 - For example: $P(A \cup B) = P(A \cap B) + P(A \cap B') + P(A' \cap B)$
 - **subtract** the probabilities of the combined outcomes that are not part of that event from 1
 - For example: $P(A \cup B) = 1 - P(A' \cap B')$



Do I have to use a tree diagram?

- If there are **multiple events** or trials then a tree diagram can get big
- You can break down the problem by using the words **AND/OR/NOT** to help you find probabilities without a tree
- You can speed up the process by only drawing parts of the tree that you are interested in

Which events do I put on the first branch?

- If the events A and B are **independent** then the **order does not matter**
- If the events A and B are **not independent** then the **order does matter**
 - If you have the probability of **A given B** then put **B on the first set** of branches
 - If you have the probability of **B given A** then put **A on the first set** of branches



Exam Tip

- In an exam do not waste time drawing a full tree diagram for scenarios with lots of events unless the question asks you to
 - Only draw the parts that you are interested in

YOUR NOTES






? Worked Example

20% of people in a company wear glasses. 40% of people in the company who wear glasses are right-handed. 50% of people in the company who don't wear glasses are right-handed.

a)


Draw a tree diagram to represent the information.



b)

One of the people in the company are randomly selected, find the probability that they are right-handed.

Find options that contain R



$$P(R) = P(G \cap R) + P(G' \cap R) = 0.08 + 0.4$$

$$P(R) = 0.48$$

c)

Given that a person who is right-handed is selected at random, find the probability that they wear glasses.

$$P(G|R) = \frac{P(G \cap R)}{P(R)} = \frac{0.08}{0.48}$$

$$P(G|R) = \frac{1}{6}$$

4.4 Probability Distributions

4.4.1 Discrete Probability Distributions

YOUR NOTES




Discrete Probability Distributions

What is a discrete random variable?

- A **random variable** is a variable whose value depends on the outcome of a **random event**
 - The value of the random variable is not known until the event is carried out (this is what is meant by 'random' in this case)
- **Random variables** are denoted using **upper case letters** (X , Y , etc)
- **Particular outcomes** of the event are denoted using **lower case letters** (x , y , etc)
- $P(X = x)$ means "the probability of the random variable X taking the value x "
- A **discrete** random variable (often abbreviated to DRV) can only take **certain values** within a set
 - Discrete random variables **usually count** something
 - Discrete random variables usually can only take a finite number of values but it is possible that it can take an infinite number of values (see the examples below)
- **Examples** of discrete random variables include:
 - The number of times a coin lands on heads when flipped 20 times
 - this has a finite number of outcomes: $\{0, 1, 2, \dots, 20\}$
 - The number of emails a manager receives within an hour
 - this has an infinite number of outcomes: $\{1, 2, 3, \dots\}$
 - The number of times a dice is rolled until it lands on a 6
 - this has an infinite number of outcomes: $\{1, 2, 3, \dots\}$
 - The number that a dice lands on when rolled once
 - this has a finite number of outcomes: $\{1, 2, 3, 4, 5, 6\}$


What is a probability distribution of a discrete random variable?

- A **discrete probability distribution** fully describes **all the values** that a discrete random variable can take along with their **associated probabilities**
 - This can be given in a **table**
 - Or it can be given as a **function** (called a discrete probability distribution function or "pdf")
 - They can be represented by **vertical line graphs** (the possible values for along the horizontal axis and the probability on the vertical axis)
- The **sum of the probabilities** of **all the values** of a discrete random variable is **1**
 - This is usually written $\sum P(X = x) = 1$
- A **discrete uniform distribution** is one where the random variable takes a finite number of values each with an **equal probability**
 - If there are n values then the probability of each one is $\frac{1}{n}$



LET x BE THE NUMBER THAT THE SPINNER LANDS ON

x	-2	0	$\frac{1}{3}$	5
$P(X=x)$	$\frac{1}{4}$	$\frac{1}{8}$	$\frac{1}{8}$	$\frac{1}{2}$

$$P(X=x) = \begin{cases} \frac{1}{8} & x = 0, \frac{1}{3} \\ \frac{1}{4} & x = -2 \\ \frac{1}{2} & x = 5 \\ 0 & \text{OTHERWISE} \end{cases}$$


How do I calculate probabilities using a discrete probability distribution?

- First **draw a table** to represent the probability distribution
 - If it is given as a function then find each probability
 - If any probabilities are unknown then use algebra to represent them
- **Form an equation** using $\sum P(X=x) = 1$
 - Add together all the probabilities and make the sum equal to 1
- To find $P(X=k)$
 - If k is a possible value of the random variable X then $P(X=k)$ will be given in the table
 - If k is not a possible value then $P(X=k) = 0$
- To find $P(X \leq k)$
 - Identify all possible values, x_i , that X can take which satisfy $x_i \leq k$
 - Add together all their corresponding probabilities
 - $P(X \leq k) = \sum_{x_i \leq k} P(X=x_i)$
 - Some mathematicians use the notation $F(x)$ to represent the cumulative distribution
 - $F(x) = P(X \leq x)$
- Using a similar method you can find $P(X < k)$, $P(X > k)$ and $P(X \geq k)$
- As all the probabilities add up to 1 you can form the following equivalent equations:
 - $P(X < k) + P(X = k) + P(X > k) = 1$
 - $P(X > k) = 1 - P(X \leq k)$
 - $P(X \geq k) = 1 - P(X < k)$

How do I know which inequality to use?

- $P(X \leq k)$ would be used for phrases such as:
 - At most, no greater than, etc
- $P(X < k)$ would be used for phrases such as:

YOUR NOTES





- Fewer than
- $P(X \geq k)$ would be used for phrases such as:
 - At least, no fewer than, etc
- $P(X > k)$ would be used for phrases such as:
 - Greater than, etc

? Worked Example

The probability distribution of the discrete random variable X is given by the function

$$P(X=x) = \begin{cases} kx^2 & x = -3, -1, 2, 4 \\ 0 & \text{otherwise.} \end{cases}$$

a)

Show that $k = \frac{1}{30}$.

Construct a table

x	-3	-1	2	4
$P(X=x)$	$9k$	k	$4k$	$16k$

Substitute in the values of x
e.g. $P(X=-3) = k(-3)^2 = 9k$

The probabilities add up to 1

$$9k + k + 4k + 16k = 1$$

$$30k = 1$$

$$k = \frac{1}{30}$$

b)

Calculate $P(X \leq 3)$.

Substitute k into the probabilities

x	-3	-1	2	4
$P(X=x)$	$\frac{3}{10}$	$\frac{1}{30}$	$\frac{2}{15}$	$\frac{8}{15}$

$$X \leq 3 : X = -3, -1, 2$$

$$P(X \leq 3) = P(X=-3) + P(X=-1) + P(X=2)$$

$$= \frac{3}{10} + \frac{1}{30} + \frac{2}{15}$$

$$P(X \leq 3) = \frac{7}{15}$$

4.4.2 Expected Values

YOUR NOTES



Expected Values $E(X)$

What does $E(X)$ mean and how do I calculate $E(X)$?

- $E(X)$ means the **expected value** or the **mean** of a **random variable X**
 - The expected value does not need to be an obtainable value of X
 - For example: the expected value number of times a coin will land on tails when flipped 5 times is 2.5
- For a **discrete** random variable, it is calculated by:
 - **Multiplying each value** of X with its corresponding **probability**
 - **Adding** all these terms together

$$E(X) = \sum xP(X = x)$$

- This is given in the **formula booklet**
- Look out for **symmetrical** distributions (where the values of X are symmetrical and their probabilities are symmetrical) as the mean of these is the same as the median
 - For example: if X can take the values 1, 5, 9 with probabilities 0.3, 0.4, 0.3 respectively then by symmetry the mean would be 5

How can I decide if a game is fair?

- Let X be the random variable that represents the **gain/loss** of a player in a game
 - X will be **negative** if there is a **loss**
- Normally the expected gain or loss is calculated by **subtracting** the **cost to play** the game from the **expected value** of the **prize**
- If $E(X)$ is **positive** then it means the player can **expect to make a gain**
- If $E(X)$ is **negative** then it means the player can **expect to make a loss**
- The game is called **fair** if the **expected gain is 0**
 - $E(X) = 0$



Worked Example

Daphne pays \$5 to play a game where she wins a prize of \$1, \$5, \$10 or \$100. The random variable W represents the amount she wins and has the probability distribution shown in the following table:

w	1	5	10	100
$P(W=w)$	0.35	0.5	0.05	0.01

a)

Calculate the expected value of Daphne's prize.

Formula booklet

Expected value of a discrete random variable X

$$E(X) = \sum x P(X=x)$$

$$E(W) = \sum w P(W=w)$$

$$= 1 \times 0.35 + 5 \times 0.5 + 10 \times 0.05 + 100 \times 0.01$$

$$\text{Expected value} = \$4.35$$

b)

Determine whether the game is fair.

A game is fair if expected gain/loss is 0

Prize - cost

$$4.35 - 5 = -0.65$$

Expected loss is \$0.65 so game is not fair

4.5 Binomial Distribution

4.5.1 The Binomial Distribution

YOUR NOTES



Properties of Binomial Distribution

What is a binomial distribution?

- A binomial distribution is a **discrete probability distribution**
- A **discrete random variable** X follows a **binomial distribution** if it **counts the number of successes** when an experiment satisfies the following conditions:
 - There are a **fixed finite number of trials** (n)
 - The outcome of each trial is **independent** of the outcomes of the other trials
 - There are **exactly two outcomes** of each trial (**success or failure**)
 - The **probability of success is constant** (p)
- If X follows a binomial distribution then it is denoted $X \sim B(n, p)$
 - n is the **number of trials**
 - p is the **probability of success**
- The **probability of failure is $1 - p$** which is sometimes denoted as q
- The formula for the probability of r **successful trials** is given by:
 - $P(X = r) = {}^n C_r \times p^r (1 - p)^{n - r}$ for $r = 0, 1, 2, \dots, n$
 - ${}^n C_r = \frac{n!}{r!(n - r)!}$ where $n! = n \times (n - 1) \times (n - 2) \times \dots \times 3 \times 2 \times 1$
 - You will be expected to use the distribution function on your **GDC to calculate probabilities** with the binomial distribution

What are the important properties of a binomial distribution?




- The **expected number (mean)** of successful trials is

$$E(X) = np$$

- You are given this in the **formula booklet**
- The **variance** of the number of successful trials is

$$\text{Var}(X) = np(1 - p)$$

- You are given this in the **formula booklet**
- Square root to get the **standard deviation**
- The distribution can be represented visually using a vertical line graph
 - If p is **close to 0** then the graph has a **tail to the right**
 - If p is **close to 1** then the graph has a **tail to the left**
 - If p is **close to 0.5** then the graph is **roughly symmetrical**
 - **If $p = 0.5$** then the graph is **symmetrical**



Modelling with Binomial Distribution

YOUR NOTES



How do I set up a binomial model?

- **Identify** what a **trial** is in the scenario
 - For example: rolling a dice, flipping a coin, checking hair colour
- **Identify** what the **successful outcome** is in the scenario
 - For example: rolling a 6, landing on tails, having black hair
- **Identify** the **parameters**
 - n is the number of trials and p is the probability of success in each trial
- Make sure you **clearly state** what your **random variable** is
 - For example, let be the number of students in a class of 30 with black hair

What can be modelled using a binomial distribution?

- Anything that satisfies the **four conditions**
- For example: let T be the number of times a fair coin lands on tails when flipped 20 times:
 - A trial is flipping a coin: There are 20 trials so **$n = 20$**
 - We can assume each coin flip does not affect subsequent coin flips: they are **independent**
 - A success is when the coin lands on tails: **Two outcomes** – tails or not tails (heads)
 - The coin is fair: The probability of tails is constant with **$p = 0.5$**
- Sometimes it might **seem like there are more than two outcomes**
 - For example: let Y be the number of yellow cars that are in a car park full of 100 cars
 - Although there are more than two possible colours of cars, here the trial is whether a car is yellow so there are two outcomes (yellow or not yellow)
 - Y would still need to fulfil the other conditions in order to follow a binomial distribution
- Sometimes a **sample may be taken from a population**
 - For example: 30% of people in a city have blue eyes, a sample of 30 people from the city is taken and X is the number of them with blue eyes
 - As long as the population is large and the sample is random then it can be assumed that each person has a 30% chance of having blue eyes

What can not be modelled using a binomial distribution?

- Anything where the number of trials is **not fixed** or is **infinite**
 - The number of emails received in an hour
 - The number of times a coin is flipped until it lands on heads
- Anything where the outcome of **one trial affects** the outcome of the **other trials**
 - The number of caramels that a person eats when they eat 5 sweets from a bag containing 6 caramels and 4 marshmallows
 - If you eat a caramel for your first sweet then there are less caramels left in the bag when you choose your second sweet
 - Anything where there are **more than two outcomes** of a trial
 - A person's shoe size
 - The number a dice lands on when rolled
 - Anything where the **probability of success changes**
 - The number of times that a person can swim a length of a swimming pool in under a minute when swimming 50 lengths

- The probability of swimming a lap in under a minute will decrease as the person gets tired
- The probability is **not constant**



Exam Tip

- An exam question might involve different types of distributions so make it clear which distribution is being used for each variable

YOUR NOTES





? Worked Example

It is known that 8% of a large population are immune to a particular virus. Mark takes a sample of 50 people from this population. Mark uses a binomial model for the number of people in his sample that are immune to the virus.

a)

State the distribution that Mark uses.

A trial is checking if a person is immune to the virus

A success is if the person is immune.

Let X be the number of people in the sample immune to the virus

$$X \sim B(50, 0.08)$$

Number of people in sample Probability of being immune to the virus

b)

State two assumptions that Mark must make in order to use a binomial model.

Mark needs to assume that:

- each person in the population has an 8% chance of being immune
- the sample is random and the people are independent
a person being immune does not affect the immunity of others

For example:
If all 50 came from the same family then they would not be independent

c)

Calculate the expected number of people in the sample that are immune to the virus.

Formula booklet

$$E(X) = 50 \times 0.08$$

$$4 \text{ people}$$

Binomial distribution $X \sim B(n, p)$	
Mean	$E(X) = np$

4.5.2 Calculating Binomial Probabilities

YOUR NOTES



Calculating Binomial Probabilities

Throughout this section we will use the random variable $X \sim B(n, p)$. For binomial, the probability of X taking a non-integer or negative value is always zero. Therefore any values of X mentioned in this section will be assumed to be non-negative integers.

How do I calculate $P(X = x)$: the probability of a single value for a binomial distribution?

- You should have a **GDC** that can calculate **binomial probabilities**
- You want to use the "**Binomial Probability Distribution**" function
 - This is sometimes shortened to BPD, Binomial PD or Binomial Pdf
- You will need to enter:
 - The 'x' value - the value of x for which you want to find $P(X = x)$
 - The 'n' value - the **number of trials**
 - The 'p' value - the **probability of success**
- Some calculators will give you the option of **listing the probabilities for multiple values of x at once**
- There is a formula that you can use but you are expected to be able to use the distribution function on your GDC
 - $P(X = x) = {}^n C_x \times p^x (1 - p)^{n - x}$

$$\blacksquare {}^n C_x = \frac{n!}{x!(n - x)!}$$

How do I calculate $P(a \leq X \leq b)$: the cumulative probabilities for a binomial distribution?

- You should have a **GDC** that can calculate **cumulative binomial probabilities**
 - Most calculators will find $P(a \leq X \leq b)$
 - Some calculators can only find $P(X \leq b)$
 - The identities below will help in this case
- You should use the "**Binomial Cumulative Distribution**" function
 - This is sometimes shortened to BCD, Binomial CD or Binomial Cdf
- You will need to enter:
 - The lower value - this is the **value a**
 - This can be zero in the case $P(X \leq b)$
 - The upper value - this is the **value b**
 - This can be n in the case $P(X \geq a)$
 - The 'n' value - the **number of trials**
 - The 'p' value - the **probability of success**

How do I find probabilities if my GDC only calculates $P(X \leq x)$?

- To calculate $P(X \leq x)$ just enter x into the cumulative distribution function
- To calculate $P(X < x)$ use:
 - $P(X < x) = P(X \leq x - 1)$ which works when X is a binomial random variable
 - $P(X < 5) = P(X \leq 4)$



- To calculate $P(X > x)$ use:
 - $P(X > x) = 1 - P(X \leq x)$ which works for any random variable X
 - $P(X > 5) = 1 - P(X \leq 5)$
- To calculate $P(X \geq x)$ use:
 - $P(X \geq x) = 1 - P(X \leq x - 1)$ which works when X is a binomial random variable
 - $P(X \geq 5) = 1 - P(X \leq 4)$
- To calculate $P(a \leq X \leq b)$ use:
 - $P(a \leq X \leq b) = P(X \leq b) - P(X \leq a - 1)$ which works when X is a binomial random variable
 - $P(5 \leq X \leq 9) = P(X \leq 9) - P(X \leq 4)$

What if an inequality does not have the equals sign (strict inequality)?

- For a binomial distribution (as it is discrete) you could **rewrite all strict inequalities** ($<$ and $>$) as **weak inequalities** (\leq and \geq) by using the identities for a binomial distribution
 - $P(X < x) = P(X \leq x - 1)$ and $P(X > x) = P(X \geq x + 1)$
 - For example: $P(X < 5) = P(X \leq 4)$ and $P(X > 5) = P(X \geq 6)$
- It helps to think about the **range of integers** you want
 - Identify the smallest and biggest integers in the range
- If your range has no minimum or maximum then use 0 or n
 - $P(X \leq b) = P(0 \leq X \leq b)$
 - $P(X \geq a) = P(a \leq X \leq n)$
- $P(a < X \leq b) = P(a + 1 \leq X \leq b)$
 - $P(5 < X \leq 9) = P(6 \leq X \leq 9)$
- $P(a \leq X < b) = P(a \leq X \leq b - 1)$
 - $P(5 \leq X < 9) = P(5 \leq X \leq 8)$
- $P(a < X < b) = P(a + 1 \leq X \leq b - 1)$
 - $P(5 < X < 9) = P(6 \leq X \leq 8)$



Exam Tip

- If the question is in context then write down the inequality as well as the final answer
 - This means you still might gain a mark even if you accidentally type the wrong numbers into your GDC



Worked Example

The random variable $X \sim B(40, 0.35)$. Find:

i)
 $P(X = 10)$.

Identify n and p $n = 40$ $p = 0.35$

Use binomial probability distribution on GDC

$$P(X = 10) = 0.057056\dots$$

$$P(X = 10) = 0.057 \text{ (3sf)}$$

ii)
 $P(X \leq 10)$.

Identify upper and lower values

$$P(X \leq 10) = P(0 \leq X \leq 10)$$

Use binomial cumulative distribution on GDC

$$P(X \leq 10) = 0.121491\dots$$

$$P(X \leq 10) = 0.121 \text{ (3sf)}$$

iii)
 $P(8 < X < 15)$.

Identify upper and lower values

$$P(8 < X < 15) = P(9 \leq X \leq 14)$$

Use binomial cumulative distribution on GDC

$$P(9 \leq X \leq 14) = 0.541827\dots$$

$$P(8 < X < 15) = 0.542 \text{ (3sf)}$$

4.6 Normal Distribution

4.6.1 The Normal Distribution

YOUR NOTES



Properties of Normal Distribution

The binomial distribution is an example of a discrete probability distribution. The normal distribution is an example of a **continuous** probability distribution.

What is a continuous random variable?

- A continuous random variable (often abbreviated to CRV) is a random variable that can take **any value** within a range of infinite values
 - Continuous random variables **usually measure** something
 - For example, height, weight, time, etc

What is a continuous probability distribution?

- A continuous probability distribution is a probability distribution in which the random variable X is continuous
- The probability of X being a **particular value is always zero**
 - $P(X = k) = 0$ for any value k
 - Instead we define the **probability density function** $f(x)$ for a specific value
 - This is a function that describes the **relative likelihood** that the random variable would be close to that value
 - We talk about the **probability** of X being within a **certain range**
- A continuous probability distribution can be represented by a continuous graph (the values for X along the horizontal axis and probability **density** on the vertical axis)
- The **area under the graph** between the points $x = a$ and $x = b$ is equal to $P(a \leq X \leq b)$
 - The **total area under the graph equals 1**
- As $P(X = k) = 0$ for any value k , it does not matter if we use strict or weak inequalities
 - $P(X \leq k) = P(X < k)$ for any value k when X is a **continuous random variable**


What is a normal distribution?

- A normal distribution is a **continuous probability distribution**
- The **continuous random variable** X can follow a normal distribution if:
 - The distribution is **symmetrical**
 - The distribution is **bell-shaped**
- If X follows a normal distribution then it is denoted $X \sim N(\mu, \sigma^2)$
 - μ is the **mean**
 - σ^2 is the **variance**
 - σ is the **standard deviation**
- If the **mean** changes then the graph is **translated horizontally**
- If the **variance** changes then the graph is **stretched horizontally**
 - A **small variance** leads to a **tall** curve with a **narrow** centre
 - A **large variance** leads to a **short** curve with a **wide** centre



SAME VARIANCES
DIFFERENT MEANS


SAME MEANS
DIFFERENT VARIANCES



Copyright © Save My Exams. All Rights Reserved

What are the important properties of a normal distribution?

- The **mean** is μ
- The **variance** is σ^2
 - If you need the **standard deviation** remember to square root this
- The normal distribution is symmetrical about
 - Mean = Median = Mode = μ
- There are the results:
 - Approximately **two-thirds (68%)** of the data lies within **one standard deviation** of the mean ($\mu \pm \sigma$)
 - Approximately **95%** of the data lies within **two standard deviations** of the mean ($\mu \pm 2\sigma$)
 - Nearly **all of the data (99.7%)** lies within **three standard deviations** of the mean ($\mu \pm 3\sigma$)



Copyright © Save My Exams. All Rights Reserved



Modelling with Normal Distribution

What can be modelled using a normal distribution?

- A lot of real-life continuous variables can be modelled by a normal distribution provided that the population is large enough and that the variable is **symmetrical** with **one mode**
- For a normal distribution X can take any real value, however values far from the mean (more than 4 standard deviations away from the mean) have a probability density of **practically zero**
 - This fact allows us to model variables that are not defined for all real values such as height and weight

What can not be modelled using a normal distribution?

- Variables which have **more than one mode** or **no mode**
 - For example: the number given by a random number generator
- Variables which are **not symmetrical**
 - For example: how long a human lives for



Exam Tip

- An exam question might involve different types of distributions so make it clear which distribution is being used for each variable



Worked Example

The random variable S represents the speeds (mph) of a certain species of cheetahs when they run. The variable is modelled using $N(40, 100)$.

a)

Write down the mean and standard deviation of the running speeds of cheetahs.

$$\mu = 40 \text{ and } \sigma^2 = 100$$

↑
Square root to get standard deviation

Mean $\mu = 40$
Standard deviation $\sigma = 10$

b)

State two assumptions that have been made in order to use this model.

We assume that the distribution of the speeds is

- symmetrical
- bell-shaped

4.6.2 Calculations with Normal Distribution

YOUR NOTES



Calculating Normal Probabilities

Throughout this section we will use the random variable $X \sim N(\mu, \sigma^2)$. For X distributed normally, X can take any real number. Therefore any values mentioned in this section will be assumed to be real numbers.

How do I find probabilities using a normal distribution?

- The **area under a normal curve** between the points $x = a$ and $x = b$ is equal to the **probability** $P(a < X < b)$
 - Remember for a normal distribution you do not need to worry about whether the inequality is strict ($<$ or $>$) or weak (\leq or \geq)
 - $P(a < X < b) = P(a \leq X \leq b)$
- You will be **expected to use** distribution functions on your **GDC** to find the probabilities when working with a normal distribution

How do I calculate $P(X = x)$: the probability of a single value for a normal distribution?

- The probability of a **single value** is **always zero** for a normal distribution
 - You can picture this as the area of a single line is zero
- $P(X = x) = 0$
- Your GDC is likely to have a "**Normal Probability Density**" function
 - This is sometimes shortened to NPD, Normal PD or Normal Pdf
 - **IGNORE THIS FUNCTION** for this course!
 - This calculates the **probability density function** at a point **NOT the probability**

How do I calculate $P(a < X < b)$: the probability of a range of values for a normal distribution?

- You need a **GDC** that can calculate **cumulative normal probabilities**
- You want to use the "**Normal Cumulative Distribution**" function
 - This is sometimes shortened to NCD, Normal CD or Normal Cdf
- You will need to enter:
 - The 'lower bound' - this is the value a
 - The 'upper bound' - this is the value b
 - The ' μ ' value - this is the mean
 - The ' σ ' value - this is the standard deviation
- **Check the order carefully** as some calculators ask for standard deviation before mean
 - Remember it is the standard deviation
 - so if you have the **variance** then **square root it**
- **Always sketch** a quick diagram to visualise which area you are looking for

How do I calculate $P(X > a)$ or $P(X < b)$ for a normal distribution?

- You will still use the "**Normal Cumulative Distribution**" function
- $P(X > a)$ can be estimated using an **upper bound that is sufficiently bigger** than the **mean**
 - Using a value that is more than 4 standard deviations **bigger than the mean** is quite accurate
 - Or an easier option is just to input lots of 9's for the upper bound (**99999999... or 10^{99}**)

- $P(X < b)$ can be estimated using a **lower bound that is sufficiently smaller** than the **mean**
 - Using a value that is more than 4 standard deviations **smaller than the mean** is quite accurate
 - Or an easier option is just to input lots of 9's for the lower bound with a negative sign (**-99999999... or -10^{99}**)

Are there any useful identities?

- $P(X < \mu) = P(X > \mu) = 0.5$
- As $P(X = a) = 0$ you can use:
 - $P(X < a) + P(X > a) = 1$
 - $P(X > a) = 1 - P(X < a)$
 - $P(a < X < b) = P(X < b) - P(X < a)$
- These are useful when:
 - The mean and/or standard deviation are unknown
 - You only have a diagram
 - You are working with the **inverse distribution**



Exam Tip

- Check carefully whether you have entered the standard deviation or variance into your GDC

YOUR NOTES





Worked Example

The random variable $Y \sim N(20, 5^2)$. Calculate:

i)
 $P(Y = 20)$.

Identify μ and σ
 $\mu = 20$ $\sigma^2 = 5^2$ so $\sigma = 5$


Sketch!



$P(Y = 20) = 0$

ii)
 $P(18 \leq Y < 27)$.

Sketch!



Using GDC

Lower = 18

Upper = 27


We can use \leq or $<$

$P(18 < Y < 27) = 0.574665\dots$

0.575 (3sf)

iii)
 $P(Y > 29)$

Sketch!



Using GDC

Lower = 29

Upper = 99999

$P(Y > 29) = 0.035930\dots$

0.0359 (3sf)

No upper bound so choose a big number

Inverse Normal Distribution

Given the value of $P(X < a)$ how do I find the value of a ?

- Your **GDC** will have a function called "**Inverse Normal Distribution**"
 - Some calculators call this InvN
- Given that $P(X < a) = p$ you will need to enter:
 - The 'area' - this is the value p
 - Some calculators might ask for the 'tail' - this is the left tail as you know the area to the left of a
 - The ' μ ' value - this is the mean
 - The ' σ ' value - this is the standard deviation

Given the value of $P(X > a)$ how do I find the value of a ?

- Given $P(X > a) = p$
- Use $P(X < a) = 1 - P(X > a)$ to rewrite this as
 - $P(X < a) = 1 - p$
- Then use the **method for $P(X < a)$** to find a
- If your calculator does have the **tail option** (left, right or centre) then you can use the "Inverse Normal Distribution" function straightaway by:
 - Selecting 'right' for the tail
 - Entering the area as ' p '



Exam Tip

- Always check your **answer makes sense**
 - If $P(X < a)$ is **less than 0.5** then a should be **smaller than the mean**
 - If $P(X < a)$ is **more than 0.5** then a should be **bigger than the mean**
 - A sketch will help you see this

YOUR NOTES





Worked Example


The random variable $W \sim N(50, 36)$.

Find the value of w such that $P(W > w) = 0.175$.

Identify μ and σ

$$\mu = 50 \quad \sigma^2 = 36 \quad \text{so } \sigma = 6$$

Sketch!



$P(W > w)$ is less than 0.5
so w is bigger than the mean

Area from left is 0.825

Use Inverse Normal Distribution function on GDC

$$w = 55.6075\dots$$

$$w = 55.6 \text{ (3sf)}$$

4.7 Hypothesis Testing

4.7.1 Hypothesis Testing

YOUR NOTES



Language of Hypothesis Testing

What is a hypothesis test?

- A hypothesis test uses a **sample of data** in an experiment to test a **statement** made about the **population**
 - The statement is either about a **population parameter** or the distribution of the **population**
- The hypothesis test will look at the probability of observed outcomes happening under set conditions
- The probability found will be compared against a given **significance level** to determine whether there is **evidence to support** the statement being made

What are the key terms used in statistical hypothesis testing?

- Every hypothesis test **must** begin with a clear **null hypothesis** (what we believe to already be true) and **alternative hypothesis** (how we believe the data pattern or probability distribution might have changed)
- A **hypothesis** is an assumption that is made about a particular **population parameter** or the **distribution of the population**
 - A **population parameter** is a numerical characteristic which helps define a population
 - Such as the mean value of the population
 - The **null hypothesis** is denoted H_0 and sets out the assumed population parameter or distribution given that no change has happened
 - The **alternative hypothesis** is denoted H_1 and sets out how we think the population parameter or distribution could have changed
 - A **one-tailed test** is used for testing the distribution or testing whether the parameter has increased (or decreased)
 - A **two-tailed test** is used for testing whether the parameter has changed (either increased or decreased)
 - When a hypothesis test is carried out, the null hypothesis is **assumed to be true** and this assumption will either be **accepted** or **rejected**
 - When a null hypothesis is accepted or rejected a **statistical inference** is made
- A hypothesis test will always be carried out at an appropriate **significance level**
 - The significance level sets the **smallest probability** that an event could have occurred by chance
 - Any probability smaller than the significance level would suggest that the event is unlikely to have happened by chance
 - The **significance level** must be set **before** the hypothesis test is carried out
 - The **significance level** will usually be 1%, 5% or 10%, however it may vary

Conclusions of Hypothesis Testing

How do I decide whether to reject or accept the null hypothesis?

- A sample of the population is taken and the **test statistic** is calculated **using the observations** from the sample
 - Your GDC will calculate the test statistic for you
- To decide whether or not to reject the null hypothesis you first need either the **p-value** or the **critical region**
- The **p-value** is the probability of a value being **at least as extreme** as the test statistic, assuming that the null hypothesis is true
 - Your GDC will give you the p-value
 - If the **p-value is less than the significance level** then the **null hypothesis** would be **rejected**
- The **critical region** is the range of values of the test statistic which will lead to the **null hypothesis** being **rejected**
 - If the **test statistic** falls within the **critical region** then the **null hypothesis** would be **rejected**
- The **critical value** is the boundary of the critical region
 - It is the least extreme value that would lead to the rejection of the null hypothesis
 - The **critical value** is determined by the **significance level**
 - In your exam you will be given the critical value if it is needed

How should a conclusion be written for a hypothesis test?

- Your conclusion **must** be written in the **context** of the question
- Use the **wording in the question** to help you write your conclusion
 - If **rejecting the null hypothesis** your conclusion should state that there is **sufficient evidence** to suggest the null hypothesis is unlikely true
 - If **accepting the null hypothesis** your conclusion should state that there is **not enough evidence** to suggest null hypothesis is unlikely true
- Your conclusion **must not** be definitive
 - There is a chance that the test has led to an **incorrect conclusion**
 - The outcome is **dependent on the sample**
 - a **different sample** might lead to a **different outcome**
- The conclusion of a two-tailed test can state if there is evidence of a change
 - You should not state whether this change is an increase or decrease



Exam Tip

- Accepting the null hypothesis does **not** mean that you are saying it is true
 - You are simply saying there is not enough evidence to reject it

YOUR NOTES



4.7.2 Chi-squared Test for Independence

YOUR NOTES



Chi-Squared Test for Independence

What is a chi-squared test for independence?

- A chi-squared (χ^2) **test for independence** is a hypothesis test used to test whether **two variables are independent** of each other
 - This is sometimes called a χ^2 **two-way test**
- This is an example of a **goodness of fit** test
 - We are testing whether the data fits the model that the variables are independent
- The chi-squared (χ^2) distribution is used for this test
- You will use a **contingency table**
 - This is a two-way table that shows the **observed frequencies** for the different combinations of the two variables
 - For example: if the two variables are hair colour and eye colour then the contingency table will show the frequencies of the different combinations

What is the degree of freedom?

- The **degree of freedom** refers to the **minimum number of expected values** you need to know in order to be able to calculate them all
- The degree of freedom is denoted ν
- For a **test for independence** with an $m \times n$ contingency table
 - $\nu = (m - 1) \times (n - 1)$
 - For example: If there are 5 rows and 3 columns then you only need to know **2 of the values** in **4 of the rows** as the rest can be calculated using the totals

What are the steps for a chi-squared test for independence?

- **STEP 1: Write the hypotheses**
 - H_0 : Variable X is independent of variable Y
 - H_1 : Variable X is not independent of variable Y
 - Make sure you clearly write what the variables are and don't just call them X and Y
- **STEP 2: Calculate the degree of freedom** for the test
 - For an $m \times n$ contingency table
 - Degree of freedom is $\nu = (m - 1) \times (n - 1)$
- **STEP 3: Enter your observed frequencies** into your GDC using the option for a 2-way test
 - Enter these as a matrix
 - Your GDC will give you a matrix of the **expected values** (assuming the variables are independent)
 - Your GDC will also give you the χ^2 statistic and its p-value
 - The χ^2 statistic is denoted as χ^2_{calc}
- **STEP 4: Decide whether there is evidence to reject the null hypothesis**
 - EITHER compare the χ^2 **statistic** with the given **critical value**
 - If χ^2 statistic $>$ critical value then **reject H_0**
 - If χ^2 statistic $<$ critical value then **accept H_0**
 - OR compare the **p-value** with the given **significance level**
 - If p-value $<$ significance level then **reject H_0**
 - If p-value $>$ significance level then **accept H_0**

• **STEP 5:** Write your **conclusion**

- If you **reject H_0**
 - There is sufficient evidence to suggest that variable X is not independent of variable Y
 - Therefore this suggests they are **associated**
- If you **accept H_0**
 - There is insufficient evidence to suggest that variable X is not independent of variable Y
 - Therefore this suggests they are **independent**

How do I calculate the chi-squared statistic?

- You are **expected** to be able to use your **GDC** to calculate the χ^2 statistic by inputting the matrix of the observed frequencies
- Seeing how it is done by hand might deepen your understanding but you are **not expected** to use this method
- **STEP 1:** For each **observed frequency** O_i calculate its **expected frequency** E_i
 - Assuming the variables are independent
 - $E_i = P(X=x) \times P(Y=y) \times \text{Total}$
 - Which simplifies to $E_i = \frac{\text{Row Total} \times \text{Column Total}}{\text{Overall Total}}$
- **STEP 2:** Calculate the χ^2 statistic using the formula
 - $\chi^2_{\text{calc}} = \sum \frac{(O_i - E_i)^2}{E_i}$
 - You do not need to learn this formula as your GDC calculates it for you
- To calculate the p -value you would find the probability of a value being bigger than your χ^2 statistic using a χ^2 distribution with v degrees of freedom



Exam Tip

Note for Internal Assessments (IA)

- If you use a χ^2 test in your IA then beware that the outcome may not be accurate if:
 - Any of the **expected values are less than 5**
 - There is **only 1 degree of freedom**
 - This means it is a 2×2 contingency table
- Note that **none of these cases will occur in the exam**

YOUR NOTES





Worked Example

At a school in Paris, it is believed that favourite film genre is related to favourite subject. 500 students were asked to indicate their favourite film genre and favourite subject from a selection and the results are indicated in the table below.

	Comedy	Action	Romance	Thriller
Maths	51	52	37	55
Sports	59	63	41	33
Geography	35	31	28	15

It is decided to test this hypothesis by using a χ^2 test for independence at the 1% significance level.

The critical value is 16.812.

a)

State the null and alternative hypotheses for this test.

H_0 : Favourite subject is independent of favourite film genre
 H_1 : Favourite subject is **not** independent of favourite film genre

b)

Write down the number of degrees of freedom for this table.

$$v = (\text{rows} - 1) \times (\text{columns} - 1) = (3 - 1) \times (4 - 1)$$

$$v = 6$$

c)

Calculate the χ^2 test statistic for this data.

Type matrix into GDC
 χ^2 statistic = 12.817...

$$\chi^2_{\text{calc}} = 12.8 \text{ (3 sf)}$$

d)

Write down the conclusion to the test. Give a reason for your answer.

$$12.8 < 16.812$$

Accept H_0 as χ^2 statistic < critical value.
There is insufficient evidence to suggest that favourite subject is not independent of favourite film genre. Therefore this suggests they are independent.

YOUR NOTES



4.7.3 Goodness of Fit Test

YOUR NOTES



Chi-Squared GOF: Uniform

What is a chi-squared goodness of fit test for a given distribution?

- A chi-squared (χ^2) **goodness of fit test** is used to test data from a sample which suggests that the population has a given distribution
- This could be that:
 - the proportions of the population for different categories follows a **given ratio**
 - the population follows a **uniform distribution**
 - This means all outcomes are **equally likely**

What are the steps for a chi-squared goodness of fit test for a given distribution?

- **STEP 1:** Write the **hypotheses**
 - H_0 : Variable X can be modelled by the given distribution
 - H_1 : Variable X cannot be modelled by the given distribution
 - Make sure you clearly write what the variable is and don't just call it X
- **STEP 2:** Calculate the degree of freedom for the test
 - For k outcomes
 - Degree of freedom is $\nu = k - 1$
- **STEP 3:** Calculate the **expected frequencies**
 - Split the total frequency using the given ratio
 - For a uniform distribution: divide the total frequency N by the number of outcomes k
- **STEP 4:** Enter the **frequencies** and the **degree of freedom** into your GDC
 - Enter the observed and expected frequencies as two separate lists
 - Your GDC will then give you the χ^2 statistic and its p -value
 - The χ^2 statistic is denoted as χ^2_{calc}
- **STEP 5:** Decide **whether** there is evidence to **reject the null hypothesis**
 - EITHER compare the χ^2 **statistic** with the given **critical value**
 - If χ^2 statistic $>$ critical value then **reject H_0**
 - If χ^2 statistic $<$ critical value then **accept H_0**
 - OR compare the **p -value** with the given **significance level**
 - If p -value $<$ significance level then **reject H_0**
 - If p -value $>$ significance level then **accept H_0**
- **STEP 6:** Write your **conclusion**
 - If you **reject H_0**
 - There is sufficient evidence to suggest that variable X does not follow the given distribution
 - Therefore this suggests that the data is **not distributed as claimed**
 - If you **accept H_0**
 - There is insufficient evidence to suggest that variable X does not follow the given distribution
 - Therefore this suggests that the data is **distributed as claimed**



Worked Example

A car salesman is interested in how his sales are distributed and records his sales results over a period of six weeks. The data is shown in the table.

Week	1	2	3	4	5	6
Number of sales	15	17	11	21	14	12

A χ^2 goodness of fit test is to be performed on the data at the 5% significance level to find out whether the data fits a uniform distribution.

a)

Find the expected frequency of sales for each week if the data were uniformly distributed.

If uniformly distributed all expected frequencies are equal

$$\text{Expected frequency} = \frac{15 + 17 + 11 + 21 + 14 + 12}{6}$$

$$\text{Expected frequency} = 15$$

b)

Write down the null and alternative hypotheses.

H_0 : Number of sales can be modelled by a uniform distribution

H_1 : Number of sales can not be modelled by a uniform distribution

c)

Write down the number of degrees of freedom for this test.

$$v = 6 - 1$$

$$v = 5$$

d)

Calculate the p-value.

Type two lists into GDC

Observed 15 17 11 21 14 12

Expected 15 15 15 15 15 15

$p = 0.4933\dots$

$p = 0.493$ (3sf)

e)

State the conclusion of the test. Give a reason for your answer.

$0.493 > 0.05$

Accept H_0 as p -value $>$ significance level
There is insufficient evidence to suggest that number of sales can not be modelled by a uniform distribution. Therefore this suggests it is uniformly distributed.

YOUR NOTES



Chi-Squared GOF: Binomial

YOUR NOTES



What is a chi-squared goodness of fit test for a binomial distribution?

- A chi-squared (χ^2) **goodness of fit test** is used to test data from a sample suggesting that the population has a **binomial distribution**
 - You will be given the value of p for the binomial distribution

What are the steps for a chi-squared goodness of fit test for a binomial distribution?

- **STEP 1: Write the hypotheses**
 - H_0 : Variable X can be modelled by the binomial distribution $B(n, p)$
 - H_1 : Variable X cannot be modelled by the binomial distribution $B(n, p)$
 - Make sure you clearly write what the variable is and don't just call it X
 - State the values of n and p clearly
- **STEP 2: Calculate the degrees of freedom for the test**
 - For k outcomes
 - Degree of freedom is $\nu = k - 1$
- **STEP 3: Calculate the expected frequencies**
 - Find the probability of the outcome using the binomial distribution $P(X = x)$
 - Multiply the probability by the total frequency $P(X = x) \times N$
- **STEP 4: Enter the frequencies and the degree of freedom into your GDC**
 - Enter the observed and expected frequencies as two separate lists
 - Your GDC will then give you the χ^2 statistic and its p -value
 - The χ^2 statistic is denoted as χ^2_{calc}
- **STEP 5: Decide whether there is evidence to reject the null hypothesis**
 - EITHER compare the χ^2 statistic with the given **critical value**
 - If χ^2 statistic $>$ critical value then **reject H_0**
 - If χ^2 statistic $<$ critical value then **accept H_0**
 - OR compare the p -value with the given significance level
 - If p -value $<$ significance level then **reject H_0**
 - If p -value $>$ significance level then **accept H_0**
- **STEP 6: Write your conclusion**
 - If you **reject H_0**
 - There is sufficient evidence to suggest that variable X does not follow the binomial distribution $B(n, p)$
 - Therefore this suggests that the data **does not follow $B(n, p)$**
 - If you **accept H_0**
 - There is insufficient evidence to suggest that variable X does not follow the binomial distribution $B(n, p)$
 - Therefore this suggests that the data **follows $B(n, p)$**



Worked Example

A stage in a video game has three boss battles. 1000 people try this stage of the video game and the number of bosses defeated by each player is recorded.

Number of bosses defeated	0	1	2	3
Frequency	490	384	111	15

A χ^2 goodness of fit test at the 5% significance level is used to decide whether the number of bosses defeated can be modelled by a binomial distribution with a 20% probability of success.

a)

State the null and alternative hypotheses.

H_0 : Number of bosses defeated can be modelled by the binomial distribution $B(3, 0.2)$
 H_1 : Number of bosses defeated can not be modelled by the binomial distribution $B(3, 0.2)$

b)

Assuming the binomial distribution holds, find the expected number of people that would defeat exactly one boss.

$$\text{Let } X \sim B(3, 0.2)$$

$$\text{Using GDC } P(X=1) = 0.384$$

$$\text{Expected } 1000 \times 0.384 = 384$$

$$\text{Expected frequency of 1} = 384$$

c)

Calculate the p-value for the test.



Find the other expected frequencies

$$\text{For } 0: 1000 \times P(X=0) = 1000 \times 0.512 = 512$$

$$\text{For } 2: 1000 \times P(X=2) = 1000 \times 0.096 = 96$$

$$\text{For } 3: 1000 \times P(X=3) = 1000 \times 0.008 = 8$$

Type two lists into GDC

Observed 490 384 111 15

Expected 512 384 96 8

$$\nu = 4 - 1 = 3$$

$$p = 0.02426 \dots$$

$$p = 0.0243 \text{ (3sf)}$$

d)

State the conclusion of the test. Give a reason for your answer.

$$0.0243 < 0.05$$

Reject H_0 as p -value $<$ significance level
 There is sufficient evidence to suggest that
 the number of bosses defeated can not be
 modelled by the binomial distribution $B(3, 0.2)$.

Chi-Squared GOF: Normal

YOUR NOTES



What is a chi-squared goodness of fit test for a normal distribution?

- A chi-squared (χ^2) **goodness of fit test** is used to test data from a sample suggesting that the population has a **normal distribution**
 - You will be given the value of μ and σ for the normal distribution

What are the steps for a chi-squared goodness of fit test for a normal distribution?

- **STEP 1:** Write the **hypotheses**
 - H_0 : Variable X can be modelled by the normal distribution $N(\mu, \sigma^2)$
 - H_1 : Variable X cannot be modelled by the normal distribution $N(\mu, \sigma^2)$
 - Make sure you clearly write what the variable is and don't just call it X
 - State the values of μ and σ clearly
- **STEP 2:** Calculate the **degrees of freedom** for the test
 - For k outcomes
 - Degree of freedom is $\nu = k - 1$
- **STEP 3:** Calculate the **expected frequencies**
 - Find the probability of the outcome using the normal distribution $P(a < X < b)$
 - Beware of unbounded inequalities $P(X < b)$ or $P(X > a)$
 - Multiply the probability by the total frequency $P(a < X < b) \times N$
- **STEP 4:** Enter the **frequencies** and the **degree of freedom** into your GDC
 - Enter the observed and expected frequencies as two separate lists
 - Your GDC will then give you the χ^2 statistic and its p -value
 - The χ^2 statistic is denoted as χ^2_{calc}
- **STEP 5:** Decide whether there is **evidence to reject the null hypothesis**
 - EITHER compare the **χ^2 statistic** with the given **critical value**
 - If χ^2 statistic $>$ critical value then **reject H_0**
 - If χ^2 statistic $<$ critical value then **accept H_0**
 - OR compare the **p -value** with the given **significance level**
 - If p -value $<$ significance level then **reject H_0**
 - If p -value $>$ significance level then **accept H_0**
- **STEP 6:** Write your **conclusion**
 - If you **reject H_0**
 - There is sufficient evidence to suggest that variable X does not follow the normal distribution $N(\mu, \sigma^2)$
 - Therefore this suggests that the data **does not follow** $N(\mu, \sigma^2)$
 - If you **accept H_0**
 - There is insufficient evidence to suggest that variable X does not follow the normal distribution $N(\mu, \sigma^2)$
 - Therefore this suggests that the data **follows** $N(\mu, \sigma^2)$



Worked Example

300 marbled ducks in Quacktown are weighed and the results are shown in the table below.

Mass (g)	Frequency
$m < 470$	10
$470 \leq m < 520$	158
$520 \leq m < 570$	123
$m \geq 570$	9

A χ^2 goodness of fit test at the 10% significance level is used to decide whether the mass of a marbled duck can be modelled by a normal distribution with mean 520 g and standard deviation 30 g.

- a)
Calculate the expected frequencies, giving your answers correct to 2 decimal places.

Let $X \sim N(520, 30^2)$

300 x probability

Mass (g)	Probability	Expected frequency
$m < 470$	0.047790...	14.34
$470 \leq m < 520$	0.452209...	135.66
$520 \leq m < 570$	0.452209...	135.66
$m \geq 570$	0.047790...	14.34

- b)
Write down the null and alternative hypotheses.

H_0 : Mass of the marbled ducks can be modelled by the normal distribution $N(520, 30^2)$
 H_1 : Mass of the marbled ducks can not be modelled by the normal distribution $N(520, 30^2)$

- c)
Calculate the χ^2 statistic.

Enter the observed and expected frequencies into

GDC $\nu = 4 - 1 = 3$

χ^2 statistic = 8.162 ...

$\chi^2_{\text{calc}} = 8.16$ (3sf)

d)

Given that the critical value is 6.251, state the conclusion of the test. Give a reason for your answer.

$8.16 > 6.251$

Reject H_0 as χ^2 statistic $>$ critical value.

There is sufficient evidence to suggest that the mass of the marbled ducks can not be modelled by the normal distribution $N(520, 30^2)$.

YOUR NOTES





Two-Sample Tests

What is a t-test?

- A t-test is used to **compare the means** of **two normally distributed** populations
- In the exam the **population variance will always be unknown**

What assumptions are needed for the t-test?

- The underlying distribution for **each variable** must be **normal**
- In the exam you will need to assume the variance for the two groups are equal
 - You will need to use the **pooled two-sample t-test**

What are the steps for a pooled two-sample t-test?

- **STEP 1: Write the hypotheses**
 - $H_0: \mu_x = \mu_y$
 - Where μ_x and μ_y are the **population means**
 - Make sure you make it clear which mean corresponds to each population
 - In words this means the **two population means are equal**
 - $H_1: \mu_x < \mu_y$ or $H_1: \mu_x > \mu_y$ or $H_1: \mu_x \neq \mu_y$
 - The alternative hypothesis will depend on what is being tested (see sections for one-tailed and two-tailed tests)
- **STEP 2: Enter the data into your GDC**
 - Enter two lists of data – one for each sample
 - Choose the pooled option
 - Your GDC will then give you the p -value
- **STEP 3: Decide whether there is evidence to reject the null hypothesis**
 - Compare the **p-value** with the given **significance level**
 - If $p\text{-value} < \text{significance level}$ then **reject H_0**
 - If $p\text{-value} > \text{significance level}$ then **accept H_0**
- **STEP 4: Write your conclusion**
 - If you **reject H_0**
 - There is sufficient evidence to suggest that the population mean of X is bigger than/smaller than/different to the population mean of Y
 - This will depend on the alternative hypothesis
 - If you **accept H_0**
 - There is insufficient evidence to suggest that the population mean of X is bigger than/small than/different to the population mean of Y
 - Therefore this suggests that the **population means are equal**

One-tailed Tests

How do I perform a one-tailed t-test?

- A **one-tailed test** is used to test one of the two following cases:
 - The population mean of X is **bigger** than the population mean of Y
 - The alternative hypothesis will be: $H_1: \mu_x > \mu_y$
 - Look out for words such as increase, bigger, higher, etc
 - The population mean of X is **smaller** than the population mean of Y
 - The alternative hypothesis will be: $H_1: \mu_x < \mu_y$
 - Look out for words such as decrease, smaller, lower, etc
- If you **reject the null** hypothesis then
 - This suggests that the population mean of X is **bigger** than the population mean of Y
 - If the alternative hypothesis is $H_1: \mu_x > \mu_y$
 - This suggests that the population mean of X is **smaller** than the population mean of Y
 - If the alternative hypothesis is $H_1: \mu_x < \mu_y$

YOUR NOTES





Worked Example

The times (in minutes) for children and adults to complete a puzzle are recorded below.

Children	3.1	2.7	3.5	3.1	2.9	3.2	3.0	2.9		
Adults	3.1	3.6	3.5	3.6	2.9	3.6	3.4	3.6	3.7	3.0

The creator of the puzzle claims children are generally faster at solving the puzzle than adults. A t-test is to be performed at a 1% significance level.

a)

Write down the null and alternative hypotheses.

Let μ_c be the population mean for children's times
and μ_A be the population mean for adults' times

$$H_0: \mu_c = \mu_A$$

$$H_1: \mu_c < \mu_A$$

It is claimed that children are quicker

b)

Find the p-value for this test.

Enter the data as two lists in GDC

Use 2-sample pooled t-test

$$p = 0.007259\dots$$

$$p = 0.00726 \text{ (3sf)}$$

c)

State whether the creator's claim is supported by the test. Give a reason for your answer.

$$0.00726 < 0.01$$

Reject H_0 as p-value < significance level.
There is sufficient evidence to suggest that children are generally faster at solving the puzzle than adults. This supports the creator's claim.

Two-tailed Tests

How do I perform a two-tailed t-test?

- A **two-tailed test** is used to test the following case:
 - The population mean of X is **different** to the population mean of Y
 - The alternative hypothesis will be: $H_1: \mu_x \neq \mu_y$
 - Look out for words such as change, different, not the same, etc
- If you **reject the null** hypothesis then
 - This suggests that the population mean of X is **different** to the population mean of Y
 - You can not state which one is bigger as you were not testing for that
 - All you can conclude is that there is evidence that the means are not equal
 - To test whether a specific one is bigger you would need to use a one-tailed test

YOUR NOTES





Worked Example

In a school all students must study either French or Spanish as well as maths. 18 students in a maths class complete a test and their scores are recorded along with which language they study.

Studies French	61	82	77	80	99	69	75	71	81
Studies Spanish	74	79	83	66	95	79	82	81	85

The maths teacher wants to investigate whether the scores are different between the students studying each language. A t-test is to be performed at a 10% significance level.

a)

Write down the null and alternative hypotheses.

Let μ_F be the population mean for scores of students of French and μ_S be the population mean for scores of students of Spanish

$$\begin{array}{l} H_0: \mu_F = \mu_S \\ H_1: \mu_F \neq \mu_S \end{array} \quad \text{Testing for a difference}$$

b)

Find the p-value for this test.

Enter the data as two lists in GDC

Use 2-sample pooled t-test

$$p = 0.47391\dots$$

$$p = 0.474 \quad (3sf)$$

c)

Write down the conclusion to the test. Give a reason for your answer.

$$0.474 > 0.1$$

Accept H_0 as p-value > significance level.
There is insufficient evidence to suggest that there is a difference between the scores.



Head to [savemyexams.co.uk](https://www.savemyexams.co.uk) for more awesome resources