

# 3.1 Thermal Concepts

## Question Paper

Course	DPIB Physics
Section	3. Thermal Physics
Topic	3.1 Thermal Concepts
Difficulty	Medium

**Time allowed:** 70  
**Score:** /56  
**Percentage:** /100

### Question 1a

This question is about modelling the thermal processes involved when a person is exercising.

When cycling, a person generates thermal energy but maintains an approximately constant temperature.

(a)

Define thermal energy and temperature and distinguish between the two concepts.

[3 marks]

### Question 1b

The following model may be used to estimate the rise in temperature of a cyclist assuming no thermal energy is lost.

A closed container holds 65 kg of water, which represents the mass of the cyclist. The water is heated at a rate of 2000 W for 20 minutes. This represents the energy generation in the cyclist.

(b)

Calculate:

(i)

the thermal energy generated by the heater.

[2]

(ii)

the temperature rise of the water, assuming no energy losses. The specific heat capacity of water is  $4200 \text{ J kg}^{-1} \text{ K}^{-1}$ .

[2]

[4 marks]

### Question 1c

The temperature rise calculated in (b) would be dangerous to the cyclist.

- (c)  
Outline one mechanism, other than evaporation, by which the container in the model would transfer energy to its surroundings.

[2 marks]

### Question 1d

A further process by which energy is lost from the cyclist is the evaporation of sweat.

The percentage of generated energy lost by sweating is 40%. The specific heat of vaporization of sweat is  $2.26 \times 10^6 \text{ J kg}^{-1}$ .

- (d)  
Using the information above, and your answer to part (b) (i), estimate the mass of sweat evaporated from the cyclist.

[3 marks]

### Question 2a

This question is about water as it changes state.

Water at constant pressure boils at a constant temperature.

- (a)  
Outline, in terms of the energy of the molecules, the reason for this.

[2 marks]

### Question 2b

In an experiment to measure the specific latent heat of vaporization of water, steam at  $100^{\circ}\text{C}$  was passed into water in an insulated container.

The following data are available.

- Initial mass of water in container =  $0.260\text{ kg}$
- Final mass of water in container =  $0.278\text{ kg}$
- Initial temperature of water in container =  $20.4^{\circ}\text{C}$
- Final temperature of water in container =  $53.4^{\circ}\text{C}$
- Specific heat capacity of water =  $4.18 \times 10^3\text{ J kg}^{-1}\text{K}^{-1}$

(b)

Show that the specific latent heat of vaporization of water is about  $1.8 \times 10^6\text{ J kg}^{-1}$ .

[4 marks]

### Question 2c

The accepted value of  $L$  is greater than that given in part (b).

(c)

Explain why, other than through experimental or calculation error, this is the case.

[2 marks]

### Question 2d

The insulated container is replaced with one made of iron and the experiment is repeated with the same starting temperature and masses of steam and water.

After a period of time, the container reaches thermal equilibrium with the water at a temperature of  $30.7\text{ }^{\circ}\text{C}$ . The specific heat capacity of iron is  $447\text{ J kg}^{-1}\text{ K}^{-1}$ .

(d)

Assuming no energy is lost to the surroundings, calculate the mass of the container.

[3 marks]

### Question 3a

This question is about thermal energy transfers involved in sweating.

(a)

Distinguish between the concepts of temperature and internal energy.

[3 marks]

### Question 3b

An athlete loses  $2.4\text{ kg}$  of water through sweat whilst training for 2 hours.

(b)

Estimate the rate of energy loss by the athlete due to sweating. The specific latent heat of vaporization of water is  $2.3 \times 10^6\text{ J kg}^{-1}$ .

[2 marks]

### Question 3c

The athlete sits down to rest on an aluminium chair of mass 40 kg following her training session.

(c)

The temperature of the athlete is 37.8 °C and the temperature of the chair is 293 K. The specific heat capacity of aluminium is 900 J kg<sup>-1</sup> K<sup>-1</sup>.

(i)

Outline two properties that can be determined by the relative temperatures of the athlete and the chair.

[2]

(ii)

Calculate the amount of energy transferred to the chair in order to change its temperature to be in thermal equilibrium with the athlete.

Assume the athlete maintains a constant temperature.

[2]

[4 marks]

### Question 3d

When the sweat evaporates from the athlete it turns from a liquid to a gas.

(d)

State, in terms of molecular structure and motion, two differences between a liquid and a gas.

[2 marks]

### Question 4a

This question is about a slowly melting iceberg.

- (a)  
Distinguish the difference between liquid water and solid ice, with reference to molecular motion and energy.

[2 marks]

### Question 4b

The following data is available regarding an iceberg:

- The iceberg has a density of  $920 \text{ kg m}^{-3}$
- The temperature of the iceberg is  $-25 \text{ }^\circ\text{C}$
- The volume of the iceberg is  $78\,000 \text{ m}^3$
- The specific latent heat of fusion of ice is  $3.3 \times 10^5 \text{ J kg}^{-1}$
- The specific heat capacity of ice is  $2.1 \times 10^3 \text{ J kg}^{-1} \text{ K}^{-1}$

- (b)  
Calculate the energy required to melt the iceberg to form water at  $0^\circ\text{C}$ .

[4 marks]

**Question 4c**

The Sun supplies thermal energy to the iceberg at an average rate of  $450 \text{ W m}^{-2}$ . Assume that the iceberg has a consistent surface area of  $312 \text{ m}^2$ .

(c)

Estimate the time taken, in years, to melt the iceberg, assuming the melted water is immediately removed, and no heat is lost to the surroundings.

**[3 marks]**

**Question 4d**

In reality, there is heat transferred between the sea, which is at a temperature greater than  $0^\circ\text{C}$ , and the iceberg.

(d)


Outline what effect this will have on the rate of melting of the iceberg.

**[2 marks]**



**Question 5a**

This question is about an experiment to examine how the specific heat capacity of water varies with temperature.



- (a)  
Draw the line of best fit for the data.

[2 marks]

### Question 5b

(b)

(i)

Determine the gradient of the line at a temperature of  $70\text{ }^{\circ}\text{C}$ .

[2]

(ii)

State the unit for the quantity represented by the gradient.

[1]

**[3 marks]**

### Question 5c

(c)

(i)

Estimate the area under the curve and give the unit.

[2]

(ii)

State what the area represents.

[1]

**[3 marks]**

**Question 5d**

(d)

The experiment used water that reached a height of 40 cm in a cylindrical can of diameter 50 cm. The average density of water between 70 °C and 90 °C is  $970 \text{ kg m}^{-3}$ .

(i)

Estimate the mass of water used in this experiment.

[2]

(ii)

Hence, estimate the amount of energy transferred heating the water from 70 °C to 90 °C.

[1]

**[3 marks]**