# 2.3 Functions Toolkit

# **Question Paper**

Course	DP IB Maths
Section	2. Functions
Topic	2.3 Functions Toolkit
Difficulty	Very Hard

Time allowed: 80

Score: /65

Percentage: /100

## Question la

The functions f and g are defined such that  $f(x) = 9x - 3x^2 - 3$  and  $g(x) = -\frac{66 + 2x}{3}$ , both for  $x \in \mathbb{R}$ .

(a) Find  $(g \circ f)(x)$ , giving your answer in the form  $(g \circ f)(x) = a(x-p)(x-q)$ .

[3 marks]

#### Question 1b

(b) Hence, or otherwise, find the *x*-intercepts of the graph of  $y = (g \circ f)(x)$ .

[1 mark]

### Question 1c

Let h(x) = 1 - 2x.

(c) Find the distance between the *y*-intercept of the graph of  $y = (f \circ h)(x)$  and the positive *x*-intercept of the graph of  $y = (g \circ f)(x)$ . Your answer should be given as an exact value.

[6 marks]

# Question 2a

Let the function f be such that  $f(x) = \sqrt{5x^2 - 11x + 6.05}$ .

Given that the inverse function  $f^{-1}$  exists, and that the domain of f is as large as possible,

(a) suggest a domain for f and write down the corresponding range.

[4 marks]

## Question 2b

(b) Based on your answer to part (a), find  $f^{-1}(\sqrt{22.05})$ .

[2 marks]

## Question 3a

Let 
$$f(x) = \sqrt{-3x^2 + 8x + 16}$$
.

(a) Write down the coordinates of the *y*-intercept of the graph of y = f(x).

[2 marks]

## Question 3b

Given that f has the largest possible valid domain,

(b) find the domain and range of f.

[6 marks]

## Question 4a

Let the function f be defined by  $f(x) = (2x^2 - 5x - 12)^{-\frac{1}{2}} - k$ , where k is a constant and where f has the largest possible valid domain.

(a) Find the domain of f.

[2 marks]

### **Question 4b**

(b) Given that that  $\lim_{x\to\infty} f(x) = -7$ , find the value of k.

[1 mark]

## Question 4c


(c) Write down the equations of any vertical and/or horizontal asymptotes on the graph of y = f(x).



Head to <u>savemy exams.co.uk</u> for more awe some resources

#### Question 5a

The following diagram shows the graph of y = f(x), for a function f that has the domain  $-3 \le x \le 3$ . Point A has coordinates (-3, 2.5) and point B has coordinates (3, -2.5). The x-intercept of the function is (2, 0) as shown.




f can be written as a piecewise function, where each of the two pieces is a linear function and where the domain of the first function is  $-3 \le x \le 2$ .

(a) Write down f(x) as a piecewise function.

[4 marks]

## Question 5b



[3 marks]

#### Question 6a

Consider the function h defined by  $h(x) = -4x^2 + 24x + 8$ ,  $x \in \mathbb{R}$ .

(a) Rewrite h(x) in the form  $a(x+b)^2 + c$ , where  $a, b, c \in \mathbb{Z}$ .

[2 marks]

## Question 6b

(b) Given that  $f(x) = (x-3)^2$  and that  $(g \circ f)(x) = h(x)$ , find g(x).

## Question 7a

The functions f and g are defined such that  $f(x) = \frac{3-2x}{5}$  and g(x) = 4x - 7, both for  $x \in \mathbb{R}$ .

- (a) Giving your answers in the form y = mx + c, find
  - (i)  $(g \circ f)(x)$
  - (ii)  $(f \circ g)(x)$ .

[4 marks]

#### Question 7b

(b) Describe a single transformation that would map the graph of  $y = (g \circ f)(x)$  onto the graph of  $y = (f \circ g)(x)$ .

[2 marks]

## Question 7c

(c) Given that  $(g \circ f)^{-1}(p) = 2$ , find the value of p.

## **Question 8a**

Let the functions f and g be defined by  $f(x) = \frac{9}{4}x^2 - 1$  and  $g(x) = x^2 - 2$ , both for  $x \ge 0$ .

- (a) Find
  - (i)  $f^{-1}(x)$
  - (ii)  $g^{-1}(x)$ .

[2 marks]

# **Question 8b**

(b) Find  $(f \circ g)(x)$  in the form  $ax^4 + bx^2 + c$ .

[2 marks]

# **Question 8c**

(c) Solve the equation  $(f \circ g)(x) = 0$ .

[3 marks]

## Question 9a

A rectangle has length l = 4x and width w = x.

- (a) Find an expression for
  - (i) the perimeter of the rectangle, P, in terms of x.
  - (ii) the area of the rectangle, A, in terms of x.

[2 marks]


## **Question 9b**

(b) Show that  $P(A) = 5\sqrt{A}$ .

[2 marks]

# Question 9c

The graph of the function P, for  $0 \le A \le 4$ , is shown below.



(c) On the grid above, draw the graph of the inverse function  $P^{-1}$ .