

# 2.5 Transformations of Graphs

## Question Paper

Course	DPIB Maths
Section	2. Functions
Topic	2.5 Transformations of Graphs
Difficulty	Hard

**Time allowed:** 120  
**Score:** /93  
**Percentage:** /100

**Question 1a**

Let  $f(x) = 2(x + 4)^3$  and  $g(x) = x^3$ , for  $x \in \mathbb{R}$ .

- (a) Give a full geometric description of the two individual transformations that can be combined to obtain the graph of  $f$  from the graph of  $g$ .

[2 marks]

**Question 1b**

The graph of  $f$  is translated by the vector  $\begin{pmatrix} 2 \\ -5 \end{pmatrix}$  to give the graph of  $h$ .

Now consider the graph of  $h$  as a transformation of the graph of  $g$ . The point A on the graph of  $h$  corresponds to the point  $(2, 8)$  on the graph of  $g$ .

- (b) Find the coordinates of A.

[4 marks]

**Question 2a**

Let  $f$  and  $g$  be functions such that  $g(x) = 2f(x - 1) + 2$ , for  $x \in \mathbb{R}$ .

The transformation that maps the graph of  $f$  onto the graph of  $g$  may be represented as a combination of two simpler transformations:

a vertical stretch by a factor of  $v$ ,

followed by

a translation by the vector  $\begin{pmatrix} a \\ b \end{pmatrix}$ .

(a) Write down the values of

(i)  $v$

(ii)  $a$

(iii)  $b$ .

[3 marks]

**Question 2b**

The point  $A(3, 4)$  on the graph of  $f$  is mapped to point  $B$  on the graph of  $g$ .


(b) Find the coordinates of  $B$ .

[3 marks]

**Question 3a**

Let  $f(x) = x^2 - 9$ , for  $x \in \mathbb{R}$ .

- (a) Sketch the graph of  $y = f(x)$  on the following grid in the interval  $0 \leq x \leq 5$ . Use an appropriate scale and clearly label any intersections the graph makes with the coordinate axes.



**[3 marks]**

**Question 3b**

(b) Find  $(f \circ f)(2)$ .

[2 marks]

**Question 3c**

The function  $g$  is obtained when the graph of  $f$  is translated by the vector  $\begin{pmatrix} 2 \\ 5 \end{pmatrix}$ .


(c) Sketch the graph of  $g(x)$  on the same grid above, also for the interval  $0 \leq x \leq 5$ .  
Clearly label any intersections the graph makes with the coordinate axes and label the graph in the form  $g(x) = ax^2 + bx + c$  where  $a$ ,  $b$  and  $c$  are constants to be determined.

[5 marks]

**Question 4a**

Let  $f(x) = 1.1e^{x-1} - 4$ , for  $-2 \leq x \leq 3$ .

- (a) Sketch the graph of  $y = f(x)$  on the grid below, clearly labelling any intersections the graph makes with the coordinate axes.



[4 marks]

**Question 4b**

The graph of  $f$  is reflected in the  $x$ -axis and then translated by the vector  $\begin{pmatrix} 2 \\ -3 \end{pmatrix}$  to obtain the graph of  $y = g(x)$ .

(b) Find an expression for  $g(x)$ .

[2 marks]

**Question 5**

The function  $f$  is defined by

$$f(x) = \begin{cases} 1 + 2x & \text{if } x \leq 2 \\ x^2 - 2x + 5 & \text{if } x > 2 \end{cases}$$

The graph of the function  $g$  is obtained by applying the following transformations to the graph of  $f$ :

a translation by the vector  $\begin{pmatrix} 2 \\ 0 \end{pmatrix}$ ,

followed by


a reflection in the  $x$ -axis.

Find an expression for  $g(x)$ .

[4 marks]

**Question 6a**

The following diagram shows the graph of  $y = f(x)$ ,  $-3 \leq x \leq 6$ .



(a) Write down the value of

(i)  $f(-2)$

(ii)  $f^{-1}(1)$ .

[2 marks]



**Question 6b**

(b) Find the value of  $(f \circ f)(0)$ .

[1 mark]

**Question 6c**

(c) Given that  $g(x) = f(x + 5) - 5$ , find the domain and range of  $g$ .

[2 marks]

**Question 7a**

Let  $v(t) = 4t^2 + 64$ , where  $t > 0$ .

The graph of a function  $g$  is obtained when the graph of  $v$  is transformed by

a vertical stretch by a factor of  $\frac{1}{8}$ ,

followed by

a translation by the vector  $\begin{pmatrix} 8 \\ 3 \end{pmatrix}$ .

(a) Find  $g(t)$ , giving your answer in the form  $at^2 + bt + c$ .

[4 marks]

**Question 7b**

A particle moves along a straight line so that its velocity in  $\text{ms}^{-1}$ , at time  $t$  seconds, is given by  $g(t)$ .


(b) Find the value of  $t$  when the particle's velocity is  $11 \text{ ms}^{-1}$ .

[2 marks]

**Question 8a**

Let  $f(x) = 2x^2 - 6x$ , for  $x \in \mathbb{R}$ .

- (a) Sketch the graph of  $y = f(x)$  on the grid below, clearly labelling the vertex as well as any intersections the graph makes with the coordinate axes.



[4 marks]

**Question 8b**

The graph of a function  $g$  is obtained from the graph of  $f$  by a reflection in the  $y$ -axis, followed by a horizontal stretch with scale factor  $\frac{1}{2}$ .

(b) Find an expression for  $g(x)$ , giving your answer in the form  $g(x) = a(x - h)^2 + k$ .

[5 marks]

**Question 9a**

Let  $f(x) = 2x^2 + bx + 8$ , for  $x \in \mathbb{R}$ , where  $b \in \mathbb{Z}$ .

Given that the equation  $f(x) = 0$  has two equal roots, and that  $b < 0$ ,

(a) find the value of  $b$ .

[3 marks]

**Question 9b**

(b) Find the coordinates of the vertex of the graph of  $f$ .

[2 marks]

**Question 9c**

The graph of a function  $g$  is obtained from the graph of  $f$  by a reflection in the  $y$ -axis, followed by a horizontal stretch with scale factor 2.

(c) Find an expression for  $g(x)$  and state the coordinates of the  $y$ -intercept of the graph of  $g$ .

[3 marks]

**Question 10a**

Let  $f(x) = 2x^2 - 12x + 10$

(a) For the graph of  $f$ , find

- (i) the  $x$ -intercepts
- (ii) the  $y$ -intercept
- (iii) the coordinates of the vertex.

**[5 marks]****Question 10b**

The graph of a function  $g$  is obtained from the graph of  $f$  by a reflection in the  $x$ -axis followed by a translation by the vector  $\begin{pmatrix} 1 \\ 6 \end{pmatrix}$ .

(b) Find  $g(x)$ , giving your answer in the form  $g(x) = a(x - h)^2 + k$ .**[3 marks]**

**Question 11a**

Consider the functions  $f$  and  $g$  defined by  $f(x) = \ln x$  and  $g(x) = \ln(2x + 5)$ , where each function has the largest possible valid domain.

(a) Write down the domain of  $g$ .

[1 mark]

**Question 11b**

The graph of  $f$  can be transformed onto the graph of  $g$  by a single translation and a single stretch, both of which are parallel to one of the coordinate axes.

(b) Describe the sequence of transformations in the case where:

- (i) the translation occurs first.
- (ii) the stretch occurs first.

[4 marks]

**Question 11c**

The graph of  $f$  can be also transformed onto the graph of  $g$  by a single translation using the vector  $\begin{pmatrix} a \\ b \end{pmatrix}$ .


(c) Find the exact values of  $a$  and  $b$ .

[3 marks]



**Question 12a**

The graph of a function  $f$  is shown below. The points  $A(1, 6)$  and  $B(3, -3)$  lie on the graph and are a local maximum and a local minimum respectively. The  $x$ -axis is an asymptote to the graph.



(a) On separate sets of axes, sketch the graphs of:

(i)  $y = 2(f(x) + 3)$ .

(ii)  $y = f(2x + 3)$ .

In each case give the coordinates of the points onto which A and B are mapped, and state the equation of the asymptote.

[6 marks]

**Question 12b**

The graph of  $y = f(x)$  is stretched horizontally by a scale factor of  $k$  then translated by the vector  $\begin{pmatrix} a \\ b \end{pmatrix}$  to map it onto the graph of  $y = f(5x + 10) + 4$ .

- (b) (i) Find the values of  $a$ ,  $b$  and  $k$ .
- (ii) Find the coordinates of the points onto which A and B are mapped.

[5 marks]

**Question 13a**

Consider the function  $f$  defined by  $f(x) = 0.4e^{x+1} - 3$ ,  $-6 \leq x \leq 3$ .

(a) Find the coordinates of

(i) the  $x$ -intercept

(ii) the  $y$ -intercept

of the graph of  $y = f(x)$ .

[2 marks]

**Question 13b**

(b) Sketch the graph of  $y = f(x)$ .

[2 marks]

**Question 13c**

The graph of  $f$  is first reflected in the  $y$ -axis and then translated by the vector  $\begin{pmatrix} 2 \\ 1 \end{pmatrix}$  to obtain the graph of a function  $g$ .

(c) Find an expression for  $g(x)$ .

[2 marks]

