

11.1 Spectroscopic Identification

Question Paper

Course	DPIB Chemistry
Section	11. Measurements & Data Processes
Topic	11.1 Spectroscopic Identification
Difficulty	Medium

Time allowed: 20
Score: /10
Percentage: /100

Question 1

Which alcohol is **not** likely to have a fragment at m/e at 43 in its mass spectrum?


- A** $(\text{CH}_3)_2\text{CHCH}_2\text{OH}$
- B** $\text{CH}_3\text{CH}(\text{OH})\text{CH}_2\text{CH}_2\text{CH}_3$
- C** $\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{OH}$
- D** $\text{CH}_3\text{CH}_2\text{CH}(\text{OH})\text{CH}_3$

[1 mark]


Question 2

Which of the compounds shown below is likely to have a fragment at $m/e = 45$ in its mass spectrum?


A




B



C



D



[1 mark]

Question 3

Chlorine has two isotopes ^{35}Cl and ^{37}Cl . Assuming in the molecule $\text{C}_4\text{H}_6\text{Cl}_4$ there is only one hydrogen and one carbon isotope, how many molecular ion peaks will be seen in its mass spectrum?

- A** 5
- B** 4
- C** 3
- D** 2

[1 mark]

Question 4

Bromine exists as two isotopes ^{79}Br and ^{81}Br , which are found in almost equal abundance.

Which of the following statements is correct?

- A** ^{79}Br is more reactive than ^{81}Br
- B** The mass spectrum of $\text{C}_3\text{H}_7\text{Br}$ has two molecular ion peaks at 122 and 124
- C** The atomic radius of ^{79}Br is less than the atomic radius of ^{81}Br
- D** The first ionisation energy of ^{79}Br is less than the first ionisation energy of ^{81}Br

[1 mark]

Question 5

Which alcohol is likely to have a fragment ion at $m/e = 31$ in its mass spectrum?

- A $(\text{CH}_3)_2\text{CHCH}_2\text{OH}$
- B $\text{CH}_3\text{CH}(\text{OH})\text{CH}_2\text{CH}_2\text{CH}_3$
- C $\text{CH}_3\text{CH}_2\text{CH}_2\text{C}(\text{OH})(\text{CH}_3)_2$
- D $\text{CH}_3\text{CH}_2\text{CH}(\text{OH})\text{CH}_3$

[1 mark]

Question 6


Which pair of compounds would you expect to both have a singly charged peak at $m/e = 29$ in the mass spectrum?

- A propan-1-ol and propanal
- B propanal and propanone
- C propan-2-ol and propanal
- D propan-1-ol and propan-2-ol

[1 mark]

Question 7

The infrared spectrum of a compound is shown below.



Use the infrared absorptions, in wavenumbers, to identify the compound

bond	wavenumber range/ cm^{-1}
O-H (alcohol)	3750 – 3200
C-H (alkane)	2962 – 2853
C-H (aldehyde)	2900 – 2820 and 2775 – 2700
C=O (aldehyde or ketone)	1740 - 1680

Which compound is shown by the infrared spectrum?

- A** propan-1-ol **B** propan-2-ol **C** propanal **D** propanone

[1 mark]

Question 8

Which of the ketones listed would **not** be expected to have a peak in its mass spectrum at $m/e = 57$?

- A** hexan-3-one, $\text{CH}_3\text{CH}_2\text{CH}_2\text{COCH}_2\text{CH}_3$
- B** pentan-3-one, $\text{CH}_3\text{CH}_2\text{COCH}_2\text{CH}_3$
- C** 3-methylbutanone, $(\text{CH}_3)_2\text{CHCOCH}_3$
- D** butanone, $\text{CH}_3\text{CH}_2\text{COCH}_3$

[1 mark]

Question 9


Which of the following statements about the mass spectrum of CH_3Br is correct?

- A** There is one peak for the molecular ion with an m/e value of 44.
- B** There is one peak for the molecular ion with an m/e value of 95.
- C** The last two peaks have abundances in the ratio 3:1 and occur at m/e values of 94 and 96.
- D** The last two peaks are of equal size and occur at m/e values of 94 and 96.

[1 mark]

Question 10

Below is a ^1H NMR spectrum for an unknown organic compound. The relative areas under the peaks are labelled



Which of the following compounds could give this spectrum?

- A propan-1-ol, $\text{CH}_3\text{CH}_2\text{CH}_2\text{OH}$
- B propan-2-ol, $\text{CH}_3\text{CH}(\text{OH})\text{CH}_3$
- C methoxyethane, $\text{CH}_3\text{OCH}_2\text{CH}_3$
- D pentan-2-one, $\text{CH}_3\text{CH}_2\text{CH}_2\text{COCH}_3$

[1 mark]