

7.1 Discrete Energy & Radioactivity

Question Paper

Course	DPIB Physics
Section	7. Atomic, Nuclear & Particle Physics
Topic	7.1 Discrete Energy & Radioactivity
Difficulty	Hard

Time allowed: 50
Score: /40
Percentage: /100

Question 1a

Transitions between three energy levels in a particular atom give rise to three spectral lines. In decreasing magnitudes, these are f_1 , f_2 and f_3 .

The equation which relates f_1 , f_2 and f_3 is:

$$f_1 = f_2 + f_3$$

(a)


Explain, including through the use of a sketch, how this equation relates f_1 , f_2 and f_3 .

[3]

[3 marks]

Question 1b

A different atom has a complete line emission spectra with a ground state energy of -10.0 eV. is:



(b)

Sketch and label a diagram of the possible energy levels for the atomic line spectra shown.

[5 marks]

[5 marks]

Question 1c

(c)

Explain the significance of an electron at an energy level of 0 eV.

[3]

[3 marks]**Question 1d**

(d)

(i)

Explain the statement 'the first excitation energy of the hydrogen atom is 10.2 eV'

[1]

(ii)

The ground state of hydrogen is -13.6 eV. Calculate the speed of the slowest electron that could cause this excitation of a hydrogen atom.

[2]

[3 marks]

Question 2a

A radioactive nucleus ${}_{85}^{229}\text{X}$ undergoes a beta-minus decay followed by an alpha decay to form a daughter nucleus ${}_{Z}^{\text{A}}\text{Y}$.

(a)

Write a decay equation for this interaction and hence determine the values of A and Z.

[2]

[2 marks]

Question 2b

Thorium, ${}_{90}^{232}\text{Th}$ decays to an isotope of Radium (Ra) through a series of transformations. The particles emitted in successive transformations are:

$\alpha \beta \beta \gamma \alpha$

(b)

Determine the resulting nuclide after these successive transformations.

[3]

[3 marks]

Question 2c

Through a combination of successive alpha and beta decays, the isotope of any original nucleus can be formed.

(c)

Explain the simplest sequence of alpha and beta decays required to do this

[3]

[3 marks]

Question 2d

A nucleus of Bohrium ${}^X_{\text{Y}}\text{Bh}$ decays to Mendeleevium ${}^{255}_{101}\text{Md}$ by a sequence of three alpha particle emissions.

(d)

Determine the number of neutrons in a nucleus of ${}^X_{\text{Y}}\text{Bh}$

[2]

[2 marks]

Question 3a

The table shows some of the isotopes of phosphorus and, where they are unstable, the type of decay.

Isotope	${}^{29}_{15}\text{P}$	${}^{30}_{15}\text{P}$	${}^{31}_{15}\text{P}$	${}^{32}_{15}\text{P}$	${}^{33}_{15}\text{P}$
Type of decay	β^+	β^+	stable		β^-

(a)

State whether the isotope ${}^{32}_{15}\text{P}$ is stable or not. If not, determine, with a reason, the type of decay it experiences.

[3]

[3 marks]

Question 3b

The isotope of phosphorus ${}_{15}^{30}\text{P}$ decays into an isotope of silicon, ${}_{Z}^A\text{Si}$.

(b)
Write a decay equation for this decay, finding the values of A and Z, and explain why each emission product occurs.

[3]

[3 marks]

Question 4a


The radioactive isotope uranium-238 decays in a decay series to the stable lead-206.

The half-life of ${}_{92}^{238}\text{U}$ is 4.5×10^9 years, which is much larger than all the other half-lives of the decays in the series.

A rock sample, when formed originally, contained 6.0×10^{22} atoms of ${}_{92}^{238}\text{U}$ and no ${}_{82}^{206}\text{Pb}$ atoms. At any given time, most of the atoms are either ${}_{92}^{238}\text{U}$ or ${}_{82}^{206}\text{Pb}$ with a negligible number of atoms in other forms in the decay series.

(a)

Sketch on the axes below the variation of number of ${}_{92}^{238}\text{U}$ atoms and the number of ${}_{82}^{206}\text{Pb}$ atoms in the rock sample as they vary over a period of 1.0×10^{10} years from its formation. Label your graphs U and Pb.



[2]

[2 marks]

Question 4b

A certain time, t , after its formation, the sample contained twice as many ${}_{92}^{238}\text{U}$ atoms as ${}_{82}^{206}\text{Pb}$ atoms.

(b) Show that the number of ${}_{92}^{238}\text{U}$ atoms in the rock sample at time t was 4.0×10^{22} .

[2]

[2 marks]

Question 4c

The ratio of the number of lead nuclei N_{Pb} to the number of uranium nuclei N_U at some time t is given by:

$$\frac{N_{Pb}}{N_U} = e^{\lambda t} - 1$$

λ is the decay constant and has a value of 1.54×10^{-10} years.

(c)

Calculate the time taken (in years) for there to be twice as many ${}_{92}^{238}\text{U}$ atoms as ${}_{82}^{206}\text{Pb}$ atoms.

[2]

[2 marks]

Question 4d

Lead-214 is an unstable isotope of lead-206. It decays by emitting a β^- particle to form bismuth-214 (Bi)

Bismuth is also unstable and has two decay modes:

- Emitting an α particle to form thallium-210 (Tl) + energy
- Emitting a β particle to form polonium-214 (Po) + energy

(d)

Write decay equations for the decay chain of lead-214 to thallium-210 and to polonium-214. Comment on the nature of the energy released.

[4]

[4 marks]

