Practice Paper 1

Question Paper

Course	DP IB Chemistry
Section	Set B
Topic	Practice Paper 1
Difficulty	Medium

Time allowed: 50

Score: /40

Percentage: /100



Head to <u>savemyexams.co.uk</u> for more awesome resources

Question 1

Excess dilute hydrochloric acid is added separately to equal masses of the four metals, calcium, zinc, magnesium and strontium. Which metal will give off the largest volume of hydrogen gas in the reaction?


_		
Δ	Cal	lcium
$\overline{}$	Cai	oulli

B zinc

C magnesium

D strontium

The glass containers X and Y are connected by a closed valve.



X contains pure CO_2 gas at 25 °C and a pressure of 1 × 10⁵ Pa. Container Y has been evacuated prior to the experiment and has a volume three times bigger than container X.

During the experiment, the valve is opened, and the temperature of the whole apparatus is raised to 160 °C.

What is the final pressure in the system?

A
$$\frac{1 \times 10^5 \times 160}{4 \times 25}$$

B
$$\frac{4 \times 10^5 \times 433}{3 \times 298}$$

C
$$1 \times 10^5 \times 433$$

 3×298

D
$$\frac{1 \times 10^5 \times 433}{4 \times 298}$$


The diagram below shows the skeletal formula of phenazine.

Phenazine

What is the empirical formula of phenazine?

- $A C_6H_6N$
- **B** $C_{12}H_8N_2$
- \mathbf{C} C_6H_4N
- $D C_{12}H_{12}N_2$

The mass spectrum of element X is shown below.



Which of the following statements is correct?

- A X has a relative atomic mass between 70 and 71
- **B** The three isotopes of X are separated after being converted to anions
- C The most abundant isotope of X contains 71 neutrons
- **D** The isotope of X with mass 72 will be deflected the most

Successive ionisation energies for an element, \mathbf{Y} , are shown in the table below.

Electrons removed	lst	2nd	3rd	4th	5th
lonisation energy / kJ mol ⁻¹	736	1450	7740	10500	13600

What is the most likely formula for the ion of \mathbf{Y} ?

- A. Y+
- B. Y²⁺
- C. Y³⁺
- D. Y⁴⁺

[1 mark]

Question 6

A spectral line that would be found in the **visible** spectrum of the hydrogen emission spectrum would be

- **A** $n_1 \rightarrow n_2$
- $\textbf{B} \qquad n_2 \rightarrow n_3$
- $\mathbf{C} \qquad \mathbf{n}_3 \rightarrow \mathbf{n}_2$
- ${\bm D} \qquad n_3 \to n_{\scriptscriptstyle \infty}$



 $Head to \underline{save my exams.co.uk} for more a we some resources$

Question 7

Use a periodic table to deduce the correct information about the element tin, Sn(Z = 50)

	Number of occupied main energy levels	Number of electrons in the highest main energy level
Α	4	4
В	4	14
С	5	4
D	5	14

[1 mark]

Question 8

Which complex is likely to be colourless?

- A. $[Zn(H_2O)_6]Cl_2$
- B. $[NH_4]_2[Fe(H_2O)_6][SO_4]_2$
- $C.K_3[Co(CN)_6]$
- D. $[Ni(NH_3)_6][BF_4]_2$

[1 mark]

Question 9

Which type of bonding can be described as 'the electrostatic attraction between positive nuclei and electrons and occurs by the sharing of electrons'?

- A. Hydrogen bonding
- B. Ionic bonding
- C. Metallic bonding
- D. Covalent bonding



Head to <u>savemyexams.co.uk</u> for more awesome resources

Question 10

Which of the following are properties of transition metals?

- I. They have variable oxidation states
- II. They can behave as catalysts
- III. They can have magnetic properties
- A. I and II only
- B. I and III only
- C. II and III only
- D. I, II and III

[1 mark]

Question 11


Which is a characteristic property of sodium oxide?

- A. It turns moist litmus paper blue
- B. It turns moist litmus paper red
- C. When it dissolves in distilled water it forms a solution with pH less than 7
- D. It reacts with magnesium metal

[1 mark]

Question 12

Sulfur forms the compound S_4N_4 with nitrogen, the structure is shown below.



Which of the following statements about S_4N_4 is correct?

- A. The sulfur atom marked x has two lone pairs
- B. The sulfur atom marked y has two lone pairs
- C. Each N atom has two lone pairs
- D. The N-S-N bond angle will be larger than S=N=S bond angle

Page 8 of 24

[1 mark]

Question 13

Which combination describes the PH_4^+ ion?

	Molecular geometry	Central atom hybridisation
Α	Tetrahedral	sp ³
В	Square planar	sp ³
С	Tetrahedral	sp ²
D	Square planar	sp ²

[1 mark]

Question 14

Which enthalpy changes correctly describe the following reactions?

Reaction 1: $HNO_3(aq) + NH_3(aq) \rightarrow NH_4NO_3(aq)$

Reaction 2: $CuCO_3(s) \rightarrow CuO(s) + CO_2(g)$

Reaction 3: $S(s) + O_2(g) \rightarrow SO_2(g)$

	ΔH ^Θ _c	ΔH ^Θ _f	ΔH ^Θ _{neut}	ΔH ^Θ _r
Α	2	2	1	3
В	3	2	1	2
С	3	3	1	2
D	2	3	1	3

Bond energy calculations show the enthalpy of combustion for propene to be -1572.0 kJ mol⁻¹.

Compound	C ₃ H ₆ (g)	CO ₂ (g)	H ₂ O (I)	H ₂ O (g)
ΔH ^Θ _f /kJ mol ⁻¹	+20.0	-393.5	-285.8	-241.8

Using the enthalpy of formation data, which calculation correctly shows the percentage error between propene's enthalpy of combustion values obtained from bond energy calculations and Hess's Law calculations, assuming the bond energy calculation value is correct?

A.
$$\frac{-1572.0}{((3 \times -393.5) + (3 \times -241.8) - (20)) - 1572.0} x \ 100$$

B.
$$\frac{(3 \times -393.5) + (3 \times -241.8) - (20)}{-1572.0} x \ 100$$

C.
$$\frac{(3 \times -393.5) + (3 \times -241.8) + (20)}{-1572.0} \times 100$$

D.
$$\frac{((3 \times -393.5) + (3 \times -241.8) - (20)) - (-1572.0)}{-1572.0} \times 100$$

[1 mark]

Question 16

The ΔG^{Θ}_f values for the following substances are shown.

Substance	$\Delta G^{\Theta_f}(kJ \text{mol}^{-1})$
NH ₃ (g)	-16.4
O ₂ (g)	0
H ₂ O(g)	-228.6
NO(g)	87.6

Which of the following is the correct calculation to determine ΔG^{Θ} ?

$$4NH_3(g) + 5O_2(g) = 6H_2O(g) + 4NO(g)$$

$$A.(-228.6 + 87.6) + (-16.4)$$

B.
$$(-16.4 \times 4) - [(-228.6 \times 6) + (87.6 \times 4)]$$

$$C.[-228.6 + (87.6 \times 4)] - (-16.4 \times 4)$$

D.
$$[(-228.6 \times 6) + (87.6 \times 4)] - (-16.4 \times 4)$$

Page 10 of 24



 $Head to \underline{savemyexams.co.uk} for more a we some resources\\$

[1 mark]

Question 17

Which statements are correct for ionic compounds?

- I. Solubility in water depends on the relative magnitude of the lattice energy compared to the hydration energy
- II. Melting points of ionic compounds increase as the size of the cation increases
- III. The enthalpy of solution for calcium chloride is represented by $CaCl_2(s) \rightarrow CaCl_2(aq)$
- A. I and II only
- B. I and III only
- C. II and III only
- D. I, II and III

A student performs to reactions and measures the rate of product formation.

Reaction 1: 1.5g of solid calcium carbonate is added to 100 cm³ of 0.5 M hydrochloric acid

Reaction 2: 100 cm³ of distilled water is then added to 100 cm³ of 0.5 M hydrochloric acid then 1.5g of solid calcium carbonate is added

The rate of reaction 1 was faster than the rate of reaction 2.

Which of the following 3 hypotheses correctly describes the difference in the rate?


- 1 Adding water reduces the frequency of collisions between reactant molecules.
- 2 Adding water reduces the proportion of effective collisions between reactant molecules.
- Adding water reduces the proportion of reactant molecules possessing the activation energy.

A 1 only **B** 1 and 2 only **C** 3 only **D** 1, 2 and 3

The stoichiometry of a catalysed reaction is shown by the equation below.

$$R(g) + S(g) \Rightarrow T(g) + U(g)$$

Two experiments were carried out in which the amount of product T was measured against time. The results are shown in the diagram below.



Which changes in the conditions from experiment 1 to experiment 2 might explain the results shown?

- 1 product *U* was continuously removed from the equilibrium mixture
- 2 a different catalyst was used
- 3 less of reactant R was used
- **A** 1 only **B** 1 and 3 only **C** 2 and 3 only **D** 1, 2 and 3




Head to <u>savemy exams.co.uk</u> for more awe some resources

Question 20

A student experimentally determined the rate expression for the reaction between iodine and propanone to be:

Rate =
$$k [H^+] [C_3 H_6 O]$$

Which graph is consistent with this information?



[1 mark]

Question 21

The following information was obtained for the rate constant, k, for a reaction at 25 °C

Α	E _a	R
$2.57 \times 10^{9} \mathrm{s}^{-1}$	96.2 kJ mol ⁻¹	8.31 J K ⁻¹ mol ⁻¹

Which expression correctly represents how to calculate the rate constant, k?

A.
$$k = (2.57 \times 10^9) \times e^{(-96.2/8.31 \times 25)}$$

B.
$$k = (2.57 \times 10^9) \times e^{(-96.2/8.31 \times 298)}$$

C.
$$k = (2.57 \times 10^9) \times e^{(-96200/8.31 \times 25)}$$

D.
$$k = (2.57 \times 10^9) \times e^{(-96200/8.31 \times 298)}$$



 $Head to \underline{savemyexams.co.uk} for more a we some resources\\$

Question 22

Consider the following reversible reaction:

$$3O_2(g) = 2O_3(g)$$

What is the value of K_c when the equilibrium concentrations are $[O_2] = 4.0 \text{ mol dm}^{-3}$ and $[O_3] = 4.0 \text{ mol dm}^{-3}$?

- A. 0.25
- B. 4
- C.16
- D. 64

[1 mark]

Question 23

Which of the following equations is used when calculating the temperature, in Kelvin, at which a reaction becomes feasible if $\Delta H^{\Theta} = x$ and $\Delta S^{\Theta} = y$.

- A. $T = \frac{x}{v}$
- B.T = xy
- C.T = x + y
- D. $T = \frac{y}{x}$

[1 mark]

Question 24

In the Brønsted-Lowry theory of acids and bases, the difference between a conjugate acid and its conjugate base is the presence of which of the following?

- A. a positive charge
- B. a pair of electrons
- C. a proton
- D. a hydrogen atom



Head to <u>savemy exams.co.uk</u> for more awe some resources

Question 25

How many different types of ions can be found in acid rain	n, assuming it contains a	mixture of sulfuric,	sulfurous, r	nitric and
nitrous acids?				

- A. 4
- B. 5
- C.6
- D.7

[1 mark]

Question 26

Which of the following statements are correct for a titration between 0.10 mol dm⁻³ propanoic acid and 0.10 mol dm⁻³ potassium hydroxide?

- A. The equivalence point will be at pH 7
- B. The salt formed will hydrolyse to form an acidic salt
- C. The salt formed will be CH₃COOK
- D. At half equivalence point $[CH_3CH_2COOH(aq)] = [CH_3CH_2COO^-(aq)]$

[1 mark]

Question 27

Which of the following statements is correct?


- A. As temperature increases, the pH value of pure water decreases
- B. As temperature decreases, the pH value of pure water decreases
- C. The pH of water is unaffected by temperature
- D. Pure water is not neutral



Head to <u>savemy exams.co.uk</u> for more awe some resources

Question 28

Below is a diagram of a voltaic cell. When the cell is running, what is happening in the salt bridge?



- **A** K⁺ ions flow to the zinc half-cell and NO₃⁻ ions flow to the copper half-cell
- ${f B}$ ${f K}^{\scriptscriptstyle +}$ ions flow to the copper half-cell and ${f NO_3}^{\scriptscriptstyle -}$ ions flow to the zinc half-cell
- **C** K⁺ and NO₃⁻ ions flow to the copper half-cell
- **D** K⁺ and NO₃⁻ ions flow to the zinc half-cell

Which of the following reactions could take place at the positive electrode (cathode) in a voltaic cell?

- I. $Cu^{2+}(aq)$ to Cu(s)
- II. $Br_2(g)$ to $Br^-(aq)$
- III. Co^{3+} (aq) to Co^{2+} (aq)
- A. I and II only
- B. I and III only
- C. II and III only
- D. I, II and III

[1 mark]

Question 30

Consider these standard electrode potentials.

$$Fe^{2+}(aq) + 2e^{-} \rightleftharpoons Fe(s) E^{\Theta} = -0.45 V$$

$$Cu^{2+}(aq) + 2e^{-} \rightleftharpoons Cu(s) E^{\Theta} = +0.15 V$$

Which is the correct working to determine E^{Θ}_{cell} ?

A.
$$E^{\Theta}_{cell} = 0.15 - (-0.45)$$

B.
$$E^{\Theta}_{cell} = 0.15 + (-0.45)$$

$$C.E_{cell}^{\Theta} = (-0.45) - 0.15$$

D.
$$E^{\Theta}_{cell} = 0.15 \times (-0.45)$$



 $Head to \underline{save my exams.co.uk} for more a we some resources$

Question 31

What types of isomerism can the following molecule show?

- I. Branch-chain
- II. Positional
- III. Functional group
- A I and II only
- B I and III only
- C II and III only
- D I, II and III



Head to <u>savemy exams.co.uk</u> for more awe some resources

Question 32

Which factors affect the rate of nucleophilic substitution in halogenoalkanes?

- I. The nature of the attacking nucleophile
- II. The structure of the halogenoalkane
- III. The identity of the halogen
- A. I and II only
- B. I and III only
- C. II and III only
- D. I, II and III

[1 mark]

Question 33

Compound **Q** has the formula CH₃CH₂CO₂CH₃

What is the name of compound **Q** and how does its boiling point compare with that of butanoic acid?

	name of Q	boiling point compared to butanoic acid
Α	methyl propanoate	lower
В	propyl methanoate	lower
С	methyl propanoate	higher
D	propyl methanoate	higher

An alkene, X, undergoes electrophilic addition with hydrogen bromide to form a halogenoalkane, Y, as the minor product.

The halogenoalkane, Y, can react with sodium hydroxide in aqueous conditions to form butan-1-ol.

What is the correct identity of the alkene, X?

- A. 2-methylpropene
- B. But-1-ene
- C. But-2-ene
- D. Pent-1-ene

[1 mark]

Question 35

In the presence of an H⁺ catalyst, compound X reacts with ethanoic acid to produce the compound below.

What is the molecular formula of compound X?

A C_4H_8O **B** $C_4H_8O_2$ **C** $C_2H_6O_2$ **D** $C_2H_6O_3$



Headto <u>savemyexams.co.uk</u> for more awesome resources

Question 36

What is this molecule called?

- A. Z-2-bromo-1-chloro-2-fluoroethene
- B. E-2-bromo-1-chloro-2-fluoroethene
- C. Z-1-bromo-2-chloro-1-fluoroethene
- D. E-1-bromo-2-chloro-1-fluoroethene

[1 mark]

Question 37

The synthesis of 2-propyl propanoate can be carried out in two steps:

$$CH_3COCH_3 \rightarrow CH_3CH(OH)CH_3$$

Ш

 $CH_3CH(OH)CH_3 \rightarrow C_2H_5COOCH(CH_3)_2$

What are the reagents needed in I and II?

	I	II
Α	potassium dichromate(VI)	sulfuric acid, propanoic acid
В	sodium borohydride	sulfuric acid, propanoic acid
С	sodium borohydride	sulfuric acid, ethanoic acid
D	potassium dichromate(VI)	sulfuric acid, ethanoic acid

Which molecule has an index of hydrogen deficiency (IHD) = 1?

- A. C₆H₁₀
- B. C₂Br₂
- C. C₄H₉N
- D. C₃H₈O

[1 mark]

Question 39

The ¹H NMR spectrum of CH₃CHCl₂ shows two signals.

What is the correct assignment of splitting patterns for these signals?

	CH_3 group	CH group
Α	doublet	quartet
В	quartet	doublet
С	singlet	singlet
D	triplet	singlet

[1 mark]

Question 40

Which row correctly describes the splitting pattern observed on the ¹H NMR spectrum for each labelled hydrogen?

- A. One doublet and four triplets
- B. One triplet, one quartet and one singlet
- C. One triplet, one doublet and three singlets
- D. One triplet, one quartet and three singlets



 $Head to \underline{save my exams.co.uk} for more awe some resources$